

Environmental Impact Study

Half Moon Bay



ECOLOGICAL SERVICES

July 22, 2021
Update: June 8 2026



Ecological Services
3803 Sydenham Road
Elginburg, Ontario K0H 1M0
Phone: (613) 376-6916
E-mail: mail@ecologicalservices.ca

June 8, 2026

Michael Kerford
712223 NB Ltd
49 Second Avenue
Uxbridge ON L9P 1J9

RE: Half Moon Bay updated Environmental Impact Study (EIS)

Dear Mr. Kerford,

Please see the following update to the July 22, 2021 EIS conducted for the Half Moon Bay development. The update has incorporated ecological changes to the site since that time, and included follow up field study efforts, changes to the Endangered Species Act, changes as a result of the Parson Inc. peer review and a Sept. 19, 2025 review from Development Services of Prince Edward County (PEC), and considerations stemming from the Nov. 19 public meeting at Shire Hall.

It is our opinion that the proposed Half Moon Bay development can occur without causing a negative impact to PEC natural heritage features if the recommended mitigation measures provided in this EIS are followed.

Respectfully submitted,

A handwritten signature in dark ink, appearing to read 'Rob Snetsinger', is written over a light, circular, textured background.

Rob Snetsinger
Ecological Services

Half Moon Bay EIS - Table of Contents

1. Overview	4
2 Summary and Recommendations.....	7
3 Policy Framework.....	9
4 Methods	11
5 Site History, Land Use, Topography, and Drainage	12
6 Ecological Land Classification (ELC)	14
7 Assessment of Natural Features.....	23
7.0 Species Conservation Act (SCA) Species.....	23
7.1 Wetland/Coastal Wetland	27
7.2 Area of Natural and Scientific Interest (ANSI).....	31
7.3 Fish Habitat.....	31
7.4 Valleylands	32
7.5 Woodlands.....	33
7.6 Wildlife Habitat	34
8. References.....	39
9. Species Lists	44
10. Qualifications.....	51

1. Overview

This Environmental Impact Study (EIS) was completed on behalf of 712223 NB Ltd. who are proposing to develop 16 lots on Flatt Point next to Half Moon Bay, Prince Edward County (Figure 1), and adjunct Lot 17 (Figure 2) at 3401 County Road 13. Please note that Figure 1 is being provided for context, and that a higher resolution concept plan will be made available by The Boulevard Group planners with the planning package.

The purpose of an EIS is to determine if significant natural heritage features are present, as defined by the Provincial Planning Statement (PPS), and the Prince Edward County Official Plan (OP), and if present, to determine if that they are at risk of a negative impact from the development. This effort is based on assessment requirements of the PPS and the OP.

The above stated intent of the EIS to focus on defined natural heritage features is noteworthy because all developments, no matter where they occur, will impact the environment. However, if any impact was the condition for development approval, then no development in the province could occur. This is well understood and why the term “negative impact” has been specifically adjoined to the significant natural heritage features that are defined within the PPS and the OP.

As noted, all sites have some ecological value and all developments, no matter where they are situated, will result in some ecological displacement. With this in mind, it is the intent of sustainable development in the province to achieve a balance between development and environmental protection. As such, the mechanisms of the EIS process are only intended to focus on the defined significant natural heritage features that are listed in the PPS and the Official Plan.

The significant natural heritage features typically listed include in planning policy, species at risk (SAR), significant wetland, significant woodland, significant valleyland, ANSI's, significant wildlife habitat, and fish habitat. Guiding this EIS effort are survey/policy manuals provided by the Ministry of Natural Resources and Forestry (MNRF) and the Ministry of Environment Conservation and Parks (MECP). If significant natural heritage features are found, and it is determined that their features or functions are at risk of a negative impact, the eventual EIS process has two possible outcomes.

1. Determine that the negative impact to significant natural heritage features is too great and that it will not be possible for a development as presented in a concept plan to be consistent with the Official Plan. An example would be if a provincially significant ANSI could lose its provincially significant status because of the degree of impact.
2. Determine that the development can be consistent with the Official Plan with respect to negative impacts, which can also be achieved with mitigation or compensation measures.



Figure 1. Half Moon Bay development proposal provided to Ecological Services in 2025. The main area of development is to occur within Flatts Point and will include 16 lots. The 17th lot is an adjunct at 3401 County Road 13 and is provided in Figure 2.




Figure 2. Lots 17 concept.

2 Summary and Recommendations

Summary Conclusion: It is our opinion that the proposed Half Moon Bay development can occur without causing a negative impact to natural heritage features if the following recommended mitigation measures are followed.

Species Conservation Act (SCA):

Blanding's Turtles: Nesting Blanding's Turtles could be attracted to newly exposed ground during construction as well as gravel or sand piles that might be used in construction in proximity to Lot 17. It is recommended that turtle exclusion fencing be maintained around the entire construction area during the construction phase of Lot 17.

SAR Bats: The SCA allows the proponent to determine if there is a high level of risk to SAR. In our opinion this is not the case and SAR bat registration with the SCA is not required as long as the following mitigation measures are followed.

1. If there is a need to remove non-red cedar trees greater than 25 cm DBH (measurement assigned by MECP 2022) within the nest/roost season of April 1 to Nov. 30, then the removal of the tree is only recommended if it can be shown that it is not an active roost tree as determined by an acoustic exit survey. Otherwise, there should be no tree clearing of non-red cedar trees greater than 25 cm DBH during the nest/roost season of April 1 to Nov. 30.
2. Unlike the rest of the lots where the first 50 m (in from the lake) of vegetation is recommended to be retained with at least 80% canopy cover, it is recommended that lots 1 and 2 retain the first 100 m. This is not intended to restrict the placement of a well head. If there is a proposal by a future landowner to develop within the 50 to 100 m woodland area of Lots 1 and 2, then it is recommended that development approvals be preceded by SAR bat surveys as per MECP (2022) and the results of these efforts be used to determine if there is a need to Register the development with the SCA. The results of that registration process would have potential impacts on how development would proceed in the 50 to 100 m woodland area.

Wetland

1. It is recommended that the lots next to wetland areas (Lots 10 to 15 and 17) have an adjacent 50 m no build buffer zone (i.e., residential dwelling, septic fields, sheds). Lot 16 is not next to wetland. It is understood that the well heads will need to be closer to the wetlands.
2. It is recommended that shrub and trees next to wetlands in the prescribed 50 m zone of Lots 10 to 15 and 17) next to wetland be maintained with at least 80% canopy cover to maintain buffer functionality. The removal of dead or dying trees (e.g., ash); non-native and invasive species (e.g., European buckthorn), or trees with a low Coefficient of Conservatism (e.g., Manitoba Maple and Crack Willow) is acceptable if the 80% canopy cover threshold is maintained. If there is a desire to remove these low Coefficient of Conservatism species in the buffer zone below the 80% threshold, they should be replaced by native tree plantings.

Fish Habitat

1. It is recommended that the lake shoreline areas for lots 1 to 11 (lots 12 to 16 are not next to fish habitat) have a 50 m no build buffer zone (i.e., residential dwelling, septic fields, sheds). It is understood that the well heads will need to be closer to the lake. Due to space constraints within lot 17, a 50 m buffer zone to the lake is not feasible, and Regulation 41/24 of 30 m will likely be applied.

2. It is recommended that shrub and trees in the lake buffer zone of Lots 1 to 11 be maintained with at least 80% canopy cover to maintain buffer functionality. The removal of dead or dying trees (e.g., ash); non-native and invasive species (e.g., European buckthorn), or trees with a low Coefficient of Conservatism (e.g., Manitoba Maple and Crack Willow) is acceptable if the 80% canopy cover threshold is maintained. If there is a desire to remove these low Coefficient of Conservatism species in the buffer zone below the 80% threshold, they should be replaced primarily by native tree species. Except for a band of trees along the shoreline that are mostly Crack Willow, the potential build area of Lot 17 is not treed and therefore maintaining an 80% crown cover is not feasible.

Woodland

Proximity/Water Protection: 1. It is recommended that the shoreline/wetland/SWH areas for lots 1 to 15 have a 50 m no build buffer zone (i.e., residential dwelling, septic fields, sheds). It is understood that the well head will need to be closer to the lake. Lot 16 is not adjacent to lake or wetland. Lot 17 is adjacent to the lake, but the potential build area is not treed, although it is recommended that shoreline trees be maintained.

Proximity/Water Protection: 2. It is recommended that shrub and trees in the 50 m buffer zone be maintained with at least 80% canopy cover to maintain buffer functionality. The removal of dead or dying trees (e.g., ash); non-native and invasive species (e.g., European buckthorn), or trees with a low Coefficient of Conservatism (e.g., Manitoba Maple and Crack Willow) is acceptable if the 80% canopy cover threshold is maintained. If there is a desire to remove these low Coefficient of Conservatism species in the buffer zone below the 80% threshold, they should be replaced primarily by native tree species.

Wildlife Habitat

Amphibian Breeding (wetland): 1. It is recommended that the combined SWT2 and “w” areas have a 50 m no build buffer zone. This is not intended to restrict the placement of the well head.

Amphibian Breeding (wetland): 2. It is recommended that shrub and trees in the 50 m buffer zone be maintained with at least 80% canopy cover to maintain buffer functionality. The removal of dead or dying trees (e.g., ash); non-native and invasive species (e.g., European buckthorn), or trees with a low Coefficient of Conservatism (e.g., Manitoba Maple and Crack Willow) is acceptable if the 80% canopy cover threshold is maintained. If there is a desire to remove these low Coefficient of Conservatism species in the buffer zone below the 80% threshold, they should be replaced primarily by native tree species.

It is recommended that land clearing occur outside of the breeding bird season (April 1 to Aug. 30) to avoid a contravention of the Migratory Birds Convention Act, which includes the Wood Thrush.

3 Policy Framework

Provincial Planning Statement (PPS) 2024

Issued under the *Planning Act*, the 2024 version of the PPS requires that municipalities consider natural heritage features in assessing development proposals. Guidance on the extent of adjacent lands is provided in a Natural Heritage Reference Manual (MNR 2010). The adjacent land width for significant natural heritage features is 120 m. From the 2024 PPS:

4.1.4 Development and site alteration shall not be permitted in:

- a) significant wetlands in Ecoregions 5E, 6E and 7E1; and*
- b) significant coastal wetlands*

4.1.5 Development and site alteration shall not be permitted in:

- b) significant woodlands in Ecoregions 6E;*
- c) significant valleylands in Ecoregions 6E;*
- d) significant wildlife habitat;*
- e) significant areas of natural and scientific interest;*
- f) coastal wetlands in Ecoregion 6E that are not subject to policy 4.1.4. b*

... unless it has been demonstrated that there will be no negative impacts on the natural features or their ecological functions.

4.1.6 Development and site alteration shall not be permitted in fish habitat except in accordance with provincial and federal requirements.

4.1.7 Development and site alteration shall not be permitted in habitat of endangered species and threatened species, except in accordance with provincial and federal requirements.

4.1.8 Development and site alteration shall not be permitted on adjacent lands to the natural heritage features and areas identified in policies 5.1.4, 5.1.5, and 5.1.6 unless the ecological function of the adjacent lands has been evaluated and it has been demonstrated that there will be no negative impacts on the natural features or on their ecological functions.

POLICY: Prince Edward County Official Plan (2021)

Note: It is our understanding that the 1998 OP is to be used for review basis on this development and not the new 2021 OP. The 2021 is referenced for guidance on EIS scope/terms of references as it was used for guidance in preparing the original EIS.

Impact assessments determine whether an activity is going to have a negative impact on a significant natural heritage feature. Field work for this project was completed under the guidance of the PPS and the Official Plan (OP) of Prince Edward County. For convenience, we are providing the wording of the 2021 OP below as a guiding document for this EIS. From the OP:

5.1.9 Environmental Impact Study (EIS)

1) An Environmental Impact Study (EIS) shall be required as part of an application for all development and site alteration applications within the identified Natural Heritage System, including all of its component natural heritage features and functions and/or on adjacent lands.

3) The EIS shall demonstrate that the proposed use will have no negative impact on significant natural heritage features and areas and their associated ecological functions (i.e. hydrological, hydrogeological, etc.).

4) Where an EIS indicates that development would have a negative impact on the Natural Heritage System and/or the natural heritage feature or area, function, attributes or linkages for which the lands were identified, the application shall not be supported or approved by the Municipality.

5) The EIS shall be prepared by a qualified professional(s) and shall be subject to the approval of the Municipality and Quinte Conservation, in consultation with any other authority having jurisdiction.

6) The Municipality may determine that a scoped EIS may be permitted for minor development proposals, and development proposals in areas that have been previously disturbed and/or where previous environmental studies have been undertaken. The extent of the scoping and identification of issues will be determined by the Municipality in collaboration with Quinte Conservation and any authority having jurisdiction, prior to undertaking the study.

Environmental Impact Study (EIS) Appendix B of the Adopted OP

Prince Edward County provides a guiding document for completing EIS's that is available on the County website. <https://www.thecounty.ca/residents/services/planning/official-plan/>

This is not an exhaustive field manual, but is intended to ensure that natural heritage features described in the PPS and the OP merit investigation for negative impacts.

Other Notable EIS related Legislation

- Fisheries Act (1985)
- Conservation Authorities Act (1990) and Ontario Regulation 41/24
- Migratory Birds Convention Act (1994)
- Species Conservation Act (2026)

4 Methods

Site screening was undertaken using the MNRF (2018) and MECP (2019) screening protocol.

Habitat communities are described following the methodology outlined in the Ecological Land Classification (ELC) manual for Southern Ontario (Lee *et al.*, 1998). Plant species were used to characterize ELC community types. The Natural Heritage Reference Manual (MNR 2010), Significant Wildlife Habitat Ecoregion Criteria Schedules (MNRF 2015) and Significant Wildlife Habitat Technical Guide (MNR 2000) were used to define significant natural features. Breeding bird surveys were based on the Ontario Breeding Bird Atlas Guide for Participants (Cadman and Kopysh, 2001) and the Canadian Wildlife Service Forest Bird Monitoring Program. The bat survey methodology was based on MNR (2011), MNRF (2014), and MNRF (2015), as well as through consultation with Michelle Karam (MECP bat specialist), bat expert Toby Thorne (Toronto Zoo), and Monique Charette (MECP biologist). Snake surveys were based on SAR snake protocols provided by MNRF (2016). The MNRF also provides protocols for targeted SAR surveys, which are applied where necessary, such as the MNR (2011) Bobolink Survey Methodology the MNR (2012) Whip-poor-will Survey Methodology, and the MNRF (2015) Blanding's Turtle Methodology.

Qualifications of the field work personnel is presented in Section 10 of this report. The timing and nature of the site visits is presented in Table 1.

Table 1: Site Visit Summary			
Date of Survey	Arrival Time	Surveyors	Main Purpose of Visit
2020			
June 26	1300	Rob Snetsinger	Wetland and Vegetation assessment
July 29	1000	Rob Snetsinger	Wetland and Vegetation assessment
2021			
May 4	2000	Kurt Hennige	Marsh Monitoring
May 17	715	Kurt Hennige Kaitlyn Closs	Breeding Birds
May 19	2030	Kurt Hennige	Night Birds Marsh Monitoring
May 31	730	Kurt Hennige	Breeding Birds
June 9	2000	Kurt Hennige	Marsh Monitoring Night Birds
June 16	900	Rob Snetsinger Dale Kristensen	ELC, vegetation features, wetland assessment, road route assessment, breeding birds
June 18	230	Rob Snetsinger	All taxa
June 22	700	Kurt Hennige	Breeding Birds
June 23	1100	Rob Snetsinger	All taxa
June 28	2300	Rob Snetsinger	Night Birds
2023			
June 9	830	Rob Snetsinger	All taxa
2025			
Nov 14	1120	Rob Snetsinger	All taxa
2026			
May 27	750	Rob Snetsinger Jo Barr	All taxa
June 7	655	Rob Snetsinger Jo Barr	All taxa

5 Site History, Land Use, Topography, and Drainage

Site History: Like much of Prince Edward County, the development site has an agricultural history. The adjacent 1880 historical map labels the development property as Flatts (or Flatt) Pt. and be part of a parcel owned by “Hicks”. The relatively shallow soils would have limited its potential for productive cash crops or orchards, and it was therefore likely used for cattle pasturing or hay production in support of the extensive dairy operations that once dominated agriculture in the County. Belden (1878) notes that the County once operated 26 cheese factories.



It is not known when farm operations on this land ceased, but agricultural activity was taking place in 1954, as shown in the adjacent image.

Expansion of trees into agricultural lands within Flatts Point has primarily been undertaken by red cedar, with lesser amounts of white and green ash. Most of the ash trees on the property had succumbed to the emerald ash borer by 2026. This red cedar expansion has continued into 2026 such that all field areas are now woodland.

The relative age of the red cedar trees suggests that agricultural abandonment has occurred at different times, with the first starting over 30 years ago.

Consequently, these lands have not had a significant amount of time to develop a diverse or significant ecology. Furthermore, red cedar is noted in the literature inhibiting ecological development.



Land Use: The subject property is within the Picton Ecodistrict 6E-15, with over half in some sort of agricultural use. About 1% is devoted to settlement and about 0.5% is devoted pits and quarries. The subject lands are bordered to north and east by Lake Ontario, and to the west and south by active agricultural lands and abandoned agricultural fields that are regenerating into a thicket community.

Soils: According to the Prince Edward County Soil Survey Report No. 10, Flatts Point is underlain by two soil types. The bulk of the point is underlain by the Ameliasburg Series, a stony alkaline clay loam of varying depths up to 1 m. There is also a band of soil running across the point represented by the Elmbrook Series, a slightly acid to neutral clay soil that is stonefree.

Topography/Drainage: Flatts Point is mostly flat, gradually sloping with a gradual downward slope towards Lake Ontario from all points within the point.

Geology: Prince Edward County as a whole is underlain by the Gull River Formation of the Simcoe Group. It consists of limestones, dolomitic limestone, and dolostone, with upper layers consisting of shale, typically overlain by shallow soils.

6 Ecological Land Classification (ELC)

Ecological land classification mapping (see Figure 3) for Flatts Point where Lots 1 to 16 are proposed is based on Lee et al. (1998), where habitat fragments less than 0.5 ha. are normally lumped in with the larger overall ELC type. Changes to the earlier ELC mapping had occurred by 2026 as a result of emerald ash borer infestation and continued expansion of red cedar. The ELC terms are described further down.



Figure 3. Half Moon Bay ELC boundaries for the area involving the 16 lots on Flatts Point are outlined in red and accompanied by boxed ELC terms. The outer land edge of the development property is shown in yellow.

ELC Sites (not from Lee et al. 1998)

Cu (Cultural): A cultural site is one that is influenced more by cultural activities than those that define the Lee et al. (1998) cultural eco-types (i.e., CUW, CUT, and CUM). Cu sites are recognized as having the lowest ecological value of all site types as they experience ongoing disturbances and are either lacking in vegetation or are dominated by non-native, and/or

invasive species. The buildings and lawns of an adjacent lot within 120 m of the development area have been labeled as Cultural.

w (Wetland). There is a small patch of ash swamp at this location, which is smaller than what has been previously reported, and at approximately 0.1 ha it is too small to meet the minimum ELC patch size of 0.5 ha. It is separated by the nearby SWT2 wetland by an access road, and as such, for ELC mapping purposes it should be considered part of the surrounding FOD7-2 woodland for which it has similar features. There is an old drainage channel at this location that would keep water out of the FOD7-2 woodland.

Trail Features. There are four maintained trails present (see red in adjacent image) where observed walking, horse riding, and ATV use have occurred, with or without permission of the landowner. These trails could also be traversed with a high clearance vehicle, but we did not observe any vehicle use on the property. They are shown here to show that there is ongoing use on the property. There are lesser used trails that are not maintained and not shown in this image.



ELC Sites (based on Lee et al. 1998)

CUW1-1 (Red Cedar Cultural Woodland Type). Previously identified as a Cultural Meadow in 2021, this site has now attained enough red cedar coverage (i.e., > 35%) to now classify as a Cultural Woodland. This was anticipated in 2021 in the previously submitted EIS and if this succession process continues, it is anticipated that the CUW1-1 area will eventually become a FOC2-1 area once it surpasses 60% red cedar crown canopy coverage.

The site is approximately 1 ha. in size and was the last remaining portion of remnant farmland yet to succeed into woodland. Aside from red cedar, it contains typical forbs and meadow grasses found elsewhere in the County such as Kentucky blue grass, timothy grass, brome, oxeye daisy, and cinquefoils. As seen in the adjacent image taken in 2021 the aggressive non-native dog strangling vine is the main ground cover plant visible, and as of 2026 it is still an aggressive presence in this area.



The species list from CUW1-1 community is provided in Table 9. 44% of plants are non-native (i.e., SNA), which is above the 20-30% average for natural areas in Ontario (Oldham et al. 1995). The average Coefficient of Conservatism (CC) value is 1.9 on a scale of 0 to 10, indicating a site with low ecological sensitivity. All but one of the native plants have a provincial rank of S5 (very common). The one exception is an S4 (common).

CUT1 (Mineral Cultural Thicket). Located to the west of the development property, but projected to contain the access road. It was identified in 2021 as a CUM/CUT (Cultural Meadow/Cultural Thicket), but as of 2025 (adjacent image), the expansion of shrubs (primarily staghorn sumac) has surpassed the 25% threshold to achieve dominance. A few small isolated meadow pockets are still present, but all are well below the 0.5 ha. required for separate classification.



The site is approximately 8 hectares and based on historic satellite images, it started out as an active farm field that when abandoned converted to meadow, and then to thicket as of 2026. Some tree growth expansion is occurring and in time, it is anticipated that it will become a Cultural Woodland as part of the next stage of ecological succession. Aside from the dominant staghorn sumac, and gray dogwood (to a lesser degree) the underlying plant community is comprised of typical meadow species found elsewhere in the County such as Kentucky blue grass, brome, oxeye daisy, bindweed, and *Fabaceae* sp. Other common plants observed include the shrubs Tartarian honeysuckle and *Rubus* sp., and some trees including red cedar, white elm, and white ash.

The species list for the CUT1 community is provided in Table 10. 46 % of plants are non-native (i.e., SNA), which is above the 20-30% average for natural areas in Ontario (Oldham et al. 1995). The average Coefficient of Conservatism (CC) value is 1.8 on a scale of 0 to 10, indicating a site with low ecological sensitivity. All but one of the native plants have a provincial rank of S5 (very common). The one exception is an S4 (common).

OAO (Open Aquatic): This refers to the adjacent lake. This ELC site type is applied to aquatic areas that do not contain aquatic vegetation.

The nearshore OAO is composed of limestone pieces over top limestone pavement. This unvegetated shallow zone extends about 6 m out into the lake before it reaches a water depth of about 1 m., and there is also no submergent vegetation observed beyond that point, although from aerial imagery the bare limestone pavement is seen to stretch as much as 60 m from shore.



BBS1-2 (Willow Gravel Shrub Beach Type). The shoreline beach area is too small to show in Figure 3 for mapping purposes as per Lee et al. (2018) but is presented here for clarity. Due to wave action, lake substrate limestone pieces are periodically washed up on shore, creating the pebble character of the beach. The width of the beach varies depending on Lake Ontario water levels and can range from as little as a few meters to about 8 m. The dominant plant vegetating the shoreline is crack willow, of which all of the shoreline trees in this photo is composed.



FOC2-1 (Dry-Fresh Red Cedar Coniferous Forest Type). This approximately 4 ha site is dominated by growths of red cedar. In the more densely grown areas, the dense canopy cover limits understory growth to a few species such as barren strawberry and mosses.

In areas where there has been a more recent intrusion of red cedar into field habitat the understory still has a meadow presence of typical forbs and grasses. The non-native and invasive European buckthorn is common throughout and the non-native invasive dog strangling vine is becoming prevalent.



The species list from the FOC2-1 woodland is provided in Table 14. 52% of plants are non-native (i.e., SNA), which is well above the 20-30% average for natural areas in Ontario (Oldham et al. 1995). The average Coefficient of Conservatism (CC) value is 1.3 on a scale of 0 to 10, indicating a site with low ecological sensitivity. All but one of the native plants have a provincial rank of S5 (very common). The one exception is an S4 (common).

FOD5-5 (Dry-Fresh Sugar Maple – Hickory Deciduous Forest Type).

This is the oldest and most diverse upland site type within the development property. Sugar maple and shagbark hickory are dominant trees in the overstory of this approximately 2.6 ha type. Other trees present include red oak, ash sp., basswood, bitternut hickory, black walnut, elm, and poplars. The trees of the sub-canopy layer reflect those of the top canopy, but with the inclusion of ironwood. The shrub layers are comprised of tree saplings and the invasive European buckthorn is a common shrub present. The ground cover is dominated by the invasive non-native garlic mustard, but also includes typical forest floor forbs such as Virginia waterleaf, herb Robert, Pennsylvania sedge, and mayapple.



Many dead ash trees were observed and evidence of Emerald Ash Borer impacts were by the characteristic bore patterns left behind in the wood.

The species list from the FOD5-5 woodland is provided in Table 11. 12% of plants are non-native (i.e., SNA), which is below the 20-30% average for natural areas in Ontario (Oldham et al. 1995). The average Coefficient of Conservatism (CC) value is 3.1 on a scale of 0 to 10, indicating a site with low ecological sensitivity. All but one of the native plants have a provincial rank of S5 (very common). The one exception is an S4 (common).

FOD7 (Fresh Moist Lowland Deciduous Forest Type). The dominant ash species that used to dominate this ELC site have mostly all succumbed to the emerald ash borer, and the earlier FOD7-2 delineation has been changed to FOD7, as no dominant deciduous trees species could be discerned in the remaining canopy that included white elm, sugar maple, hickory, and red cedar. The shrub layer is heavily dominated by the non-native European buckthorn and prickly ash. The ground cover is dominated by representative tree seedlings, as well as prickly ash and buckthorn seedlings. Understory ground cover forbs are relatively sparse, with poison ivy often prevalent.

The species list from the FOD7 woodland is provided in Table 12. 22% of plants are non-native (i.e., SNA), which is within the 20-30% average for natural areas in Ontario (Oldham et al. 1995). The average Coefficient of Conservatism (CC) value is 2.6 on a scale of 0 to 10, indicating a site with low ecological sensitivity. All but one of the native plants have a provincial rank of S5 (very common). The one exception is an S4 (common).

FOM (Mixed Forest). As this site was on private property, our assessment is based on a roadside analysis. A mixed woodland must have at least 25% canopy cover each of both deciduous trees and coniferous trees. The canopy in 2021 appeared to be dominated by green ash, but these mostly appear to have succumbed to the emerald ash borer. A mix of other

deciduous species were observed. The shrub layer is relatively dense and is dominated by prickly ash and European buckthorn, indicating a more recent forest succession.

The species list from the FOM woodland is provided in Table 13. 26% of plants are non-native (i.e., SNA), which is within the 20-30% average for natural areas in Ontario (Oldham et al. 1995). The average Coefficient of Conservatism (CC) value is 2.4 on a scale of 0 to 10, indicating a site with low ecological sensitivity. All but one of the native plants have a provincial rank of S5 (very common). The one exception is an S4 (common).

FOD (Deciduous Woodland). Located on adjacent lands to the development along the edge of the access road on a high point of land overlooking the adjacent wetland. The top canopy is dominated by Manitoba maple and the understory is dominated by weedy non-native species. It contains many of the same species as the nearby FOD7-2 woodland.

SWT2 (Mineral Thicket Swamp Ecosite). A more detailed ecotype could not be provided due to plant variability. It is part of a diffuse thicket swamp that is largely defined by Lake Ontario water levels. It is dominated by shrubs, but also contains up to 25% tree cover, primarily composed of green ash, American elm, and willow.

From our edge analysis it appears to be primarily bordered by silky dogwood, but also contains other shrub species including slender willow, black alder, red-osier dogwood, nannyberry. Ground cover included various obligate and facultative wetland plants such as water willow, sensitive fern, touch me net, wetland sedge/grasses, and water parsnip.



Lot 17 ELC

The ELC types associated with Lot 17 are provided in Figure 4.



Figure 4. ELC associated with Lot 17. ELC boundaries are outlined in red, and Lot 17 is outlined in yellow.

Cultural: Includes developed areas. These are located south and east of Lot 17 and part of the eastern and southern ends of Lot 17, which includes the access road.

Cultural Thicket (CUT): As noted by Lee et al (1998), cultural sites are those that have resulted from, or are maintained by anthropogenic based disturbances. From historic aerial photography the site was used as an orchard at one time. It is not known when orchard activities ended, but the lack of remnant orchard trees today suggests they were removed, perhaps to make way to some other use. It normally takes less than 20 years for a fallow site to succeed from field to woodland (i.e., 60% crown cover), and so the continued presence of field and shrub species is informative of recent disturbances. Recent lawnmower sized pathways had been cut through this property.



Cultural sites, such as this one, are generally considered to have a relatively low ecological value as they often contain many non-native species. The dominant shrub cover is staghorn sumac, but other shrubs present include Virginia creeper, grape, red cedar, raspberry species, and European buckthorn. Tree species are beginning to invade this thicket community, including white and green ash, sugar maple, basswood, and black walnut. As with most undeveloped shorelines in Prince Edward County, the shoreline is dominated by a narrow band of crack willow as shown in the adjacent image.



Many of the ground cover forbs are non-native weedy species such as wild carrot, horsetweed, daisy fleabane, vetch, and thistles.

Open Aquatic (OAO): This refers to the adjacent lake. This ELC site type is applied to aquatic areas that do not contain aquatic vegetation. The nearshore OAO is composed of limestone pieces over top limestone pavement. This unvegetated shallow zone extends about 6 m out into the lake before it reaches a water depth of about 1 m. There was no submergent vegetation observed beyond that point. Due to wave action, the underlying limestone pieces are periodically washed up on shore, creating the pebble character of the beach.

Dry-Fresh Sugar Maple – Ironwood Deciduous Forest Type (FOD5-4). A common deciduous forest type in the region. Sugar maple is the dominant tree in the overstory. Other trees present include ironwood, white ash, oaks, basswood, bitternut hickory, black walnut, and poplars. A shrub layer is mostly lacking, but the invasive European buckthorn is the most common shrub present. Evidence of the Emerald Ash Borer is present, and if similar trajectories of other sites are followed, most of the ash trees here will succumb to this invasive pest. There is limited sapling regeneration and shrub growth, which is typical for these woodlands at this stage of their development. Regeneration would further be inhibited by the allelopathic influence of the dense garlic mustard ground cover (see adjacent image). Other common ground cover species observed included showy tick trefoil, poison ivy, and mayapple.



Silky Dogwood Mineral Thicket Swamp Type (SWT2-8):

This ELC type is located more than 120 m northwest of the proposed build area (see Figure 2). As the name implies it is dominated by silky dogwood, although there may be different dominant species.

Water levels have a seasonal character, as was evident when comparing levels on June 26, 2020 to those on July 29, 2020. The adjacent image of the SWT2-8 wetland was taken on the 29th where its dry condition is evident.



7 Assessment of Natural Features

7.0 Species Conservation Act (SCA) Species

The following Species at Risk (SAR) relevant to the 2026 SCA are discussed because of their potential to be on or within 10 km of the site. The listing below was based on the MNR (2018) and MECP (2019) screening protocols, as well as past work we have done in the County. Species with only a historical presence (e.g., Henslow Sparrow) were not included.

Table 2. SCA applicable species.					
Species	ESA Status	Habitat	Habitat Present within Subject Lands	Known Sightings nearby including those more than 120 m or Observed by ES	SCA Registration Relevance
Plants					
Butternut	Endangered	Woodland	Yes	None	None
Black Ash	Endangered	Wetland	Yes	None	None
Four Leaved Milkweed	Endangered	Dry woodland slopes	No	None, most PEC sightings are from east of Picton (McMahon Bluff/Maccauley Mountain)	None
Herps					
Blanding's Turtle	Threatened	Aquatic Habitat	No	Historical only	See below
Birds					
Short Eared Owl	Threatened	Grasslands	SE adjacent fields	None	None
Mammals					
Eastern Small Footed Myotis	Endangered	Cliffs like habitat	No	No	None
Little Brown Myotis	Endangered	Cosmopolitan	Yes	No	See Below
Northern Myotis	Endangered	Cosmopolitan	Yes	No	See Below
Tri-colored Bat	Endangered	Cosmopolitan	Yes	No	See Below
Eastern Red	Endangered	Cosmopolitan	Yes	No	See Below
Silver Haired	Endangered	Cosmopolitan	Yes	No	See Below
Hoary	Endangered	Cosmopolitan	Yes	No	See Below
Fish					
American Eel	Endangered	Lake with deeper water and soft substrates	No	No (offshore waters lack suitable burrowing and feeding habitat)	None

Blanding's Turtles (THR). The closest core habitat area for the Blanding's Turtle is within the wetland associated with the Little Bluffs Conservation Area about 1 km to the east of the development property, and Blanding's Turtles have been observed here. This wetland and the surrounding bluffs area is a core habitat area because it provides nesting, feeding, and overwintering opportunities. Similar core Blanding's Turtle areas where they have been observed would be the South Bay Marsh significant wetland about 3 km to the southwest, South Bay Coastal significant wetland starting about 2 km to the south, Ostrander Point significant wetland about 3 km southeast, and Big Sand Bay significant wetland about 6 km to the southeast.

Core habitat features are described as follows:

Overwintering: This occurs in water that is deep enough to prevent ice from freezing to the bottom. While Blanding's Turtles can rest on the bottom surface of a water body, they will

occasionally burrow into the bottom sediments, or hide in muskrat houses, or hide under logs to prevent, among other things, predation by otters. The water depth in the swamp is determined by Lake Ontario water levels, via the permeable beach bar. During low water levels, including fall drawdown, the swamp dries up making it unsuitable for overwintering. It also has a mostly hardened substrate, so overwintering burrowing would be inhibited as well. The shallow depth of the non-vegetated limestone pavement that exists immediately offshore is also unsuitable for overwintering and these areas would be also subject to harmful ice scouring.

Feeding: Habitat features, including feeding areas, are described by Hartwig and Kiviat (2007), Wieten et al. (2012) and Whitney et al. (2014). Ideal sites include sufficient open water to provide water cover, and emergent vegetation and diverse submergent vegetation that provides macro/micro invertebrate diversity. These features are present at the five previously listed core habitat areas. In contrast, the waters off-shore of the development area are barren, lacking both sediments that could provide invertebrate food items and vegetation. The swamp habitat next to the development also has limited feeding value, as it often lacks standing water, and the heavy shading and dense woody vegetation limits its value as a productive feeding area.

Nesting: Successful Blanding's Turtle egg laying sites require sun exposure (i.e., no overhanging vegetated shading), an appropriate diggable substrate such as sand/gravel and a lack of nest predators (e.g., raccoons and skunks). It should also be in reasonable proximity to core overwintering and feeding areas. There are no appropriate nesting features within the 16 lots of the main development area. It is too heavily shaded or covered in dense ground vegetation for sun exposed nesting. The pebble beaches next to the development area are too narrow, too shaded, and too prone to wave wash to provide turtle nesting and no depredated nests were observed here during the field work. The proposed adjunct Lot 17 also lacks suitable nesting features, but the adjacent lawn and vehicle access areas of the residences to Lot 17 do provide suitable nesting features.

The bulk of a Blanding's Turtle population will spend most of their life in proximity to the core habitat area for feeding, nesting, and overwintering purposes. Some individuals will move further afield for nesting, but always within proximity to aquatic habitat during those movements. We note the Blanding's Turtle photographs provided to us that were taken near the residence of the Gadbois family residence on April 4, 2022 and June 24, 2025. This residence has frontage on County Road 13 and is located near the access road for Lot 17, the single adjunct lot east of the proposed area that is proposed to contain 16 lots. From the June photographs, and from a provided description of where the photo was taken, it would appear that the June turtle is in a suitable area for nesting purposes, and June is the nesting time. Given the propensity for nesting travel to occur within proximity to aquatic habitat (e.g., Lake Ontario in this instance), it is our opinion that potential nesting on the Gadbois family residence property would be from turtles traveling out from the Little Bluffs Conservation area.

Aside from nesting, some individuals from the core population will also range further afield to seek out new core habitats, akin to human colonial expansion. These efforts are not always successful, but if new core habitat is found, they do allow for territorial expansion. Given the unsuitability of the swamp wetland next to the development for overwintering, the April sightings on the Gadbois residence are likely turtles seeking new habitat opportunities from nearby core habitat areas, such as Little Bluffs. However, as previously discussed, the swamp habitat next to

the development is unsuitable as core habitat and so core habitat expansion would not be successful.

The 2026 SCA aims to protect the immediate area around key habitat features. This would include nesting areas, overwintering areas, and feeding areas. At certain times of the year, parts of the wetland associated with the development could provide some limited feeding opportunities, but lake water drawdown would prevent it from use for overwintering. As such, it would not be a core habitat area from which turtles would expand outwards for nesting. Nevertheless, there are no plans to develop within any aquatic or wetland habitat, and these habitats will also be protected with setbacks via Ontario Regulation 41/24 and recommended setbacks outlined in this EIS. In summary, there will be no loss of Blanding's Turtle nesting, overwintering, or feeding habitat from the proposed development and as such, no violation of the 2025 ESA.

Blanding's Turtle Recommendation: Nesting Blanding's Turtles could be attracted to newly exposed ground during construction as well as gravel or sand piles that might be used in construction in proximity to Lot 17. It is recommended that turtle exclusion fencing be maintained around the entire construction area during the construction phase of Lot 17.

Given the proximity to potential nesting to Lot 17 it is also recommended that the SCA registration process be followed prior to the development of Lot 17.

<https://www.ontario.ca/page/species-conservation-registry>

Bat SAR (END): All bat species added to the Ontario Endangered Species Act was as a result of White Nose Syndrome and not from habitat loss. SAR bat species were not habitat limited prior to the fungus appearing in the province, and are definitely not habitat limited today. Within several years this fungus has been able to decimate population numbers because it attacks bats when they hibernate, and since the SAR Bats congregate in large numbers in a limited number of hibernation sites, the fungus has the potential to spread through these congregations and wipe out whole populations. There are no SAR bat hibernation sites on the property.

Parts of the development property is comprised of red cedar dominated woodlands and there is an extensive literature (Table 3) that has shown that red cedar woodlands (see FOC2-1 and FOMa, Figure 3) are not favored for bat use. It is intended that most of the tree clearing for development purposes will occur in the red cedar woodlands.

Table 3. A sampling of red cedar associated bat research, which includes research from nearby areas indicating that red cedar use is not favored by bats. Some of these research papers focus on specific tree species usage, and most focus on tree characteristics are of a type that red cedar does not attain (i.e., height, cavities, snags).

Jung, T., I. Thompson, and R. Titman. 2004. Roost site selection by forest-dwelling male *Myotis* in central Ontario, Canada. *Forest Ecology and Management* 202:325-335.

Key point: Favorability of large old trees, a feature that red cedars do not attain.

Holt, R. 2022. Forest roost use by Little Brown Bats (*Myotis lucifugus*) in Ontario. M.Sc. Thesis. Trent University.

Key point: Specific trees and tree characteristics discussed, exclusive of red cedar.

Gerson, H. 1984. Habitat management guidelines for bats of Ontario. Ontario Ministry of Natural Resources.

Key point: Specific trees and tree characteristics discussed, exclusive of red cedar

Kalcounis-Ruppell, M., J. Psyllakis, and M. Brigham. 2005. Tree roost selection by bats: an empirical synthesis using meta-analysis. *Wildlife Society Bulletin*. 33:1123-1132.

Key point: Preference for tall large diameter trees.

Garroway, C. J. and H. G. Broders. 2008. Day roost characteristics of northern long-eared bats (*Myotis septentrionalis*) in relation to female reproductive status. *Ecoscience* 15(1): 89-93.

Key point: Preference for tall large diameter trees.

Olson, C. R. and R. M. Barclay. 2013. Concurrent changes in group size and roost use by reproductive female little brown bats (*Myotis lucifugus*). *Canadian Journal of Zoology* 91(3): 149-155.

Key point: Preference for tall large diameter trees.

Caceres, M. C. and R. M. Barclay. 2000. *Myotis septentrionalis*. *Mammalian Species* 634: 1-4.

Key point: Preference for tall large decaying trees.

Huynh, H. M. 2009. Another record of foliage roosting in the Little Brown Bat, *Myotis lucifugus*, in Canada. *The Canadian Field-Naturalist* 123(3): 265.

Key Point: Exclusive or red cedar.

Kurta, A. S. Murray, and D. Miller. 2002. Roost selection and movements across the summer landscape. Conference Paper on the Biology and Management of an Endangered Species.

Key Point: Exclusive or red cedar.

Lewis, M., G. Turner, M. Scafani, and J. Johnson. 2022. Seasonal roost selection and activity of a remnant population of northern myotis in Pennsylvania. *PLoS ONE* 17(7): e0270478.

Key Point: Exclusive or red cedar.

We have been conducting acoustic monitoring for the last nine years and have never found a site that did not have SAR bats. They would be picked up by the acoustic monitors in any part of PEC, including downtown Picton. We note that acoustic monitoring picks up bats while in flight and therefore this does not necessarily indicate roost use, and we note that the amended 2025 ESA and its replacement, the 2026 SCA, only focuses on roost sites.

The methodology used for roost considerations for this EIS is outlined in MNR (2011b) and it has since been reproduced by MECP (2022). Both methodologies set the threshold for significant snag numbers at 10/ha. If a site has this many snags, then it is considered a high value bat woodland that merits acoustic monitoring and possible further consideration. The number of snag plots required for the entire property is sixteen, but we did 20 plots, and based on MNR (2011b) plot modeling, the total snag count was 3/ha., which is well below 10/ha. Nevertheless, we do recognize the bat potential of the FOD5-5 woodland and placed bat monitors here in 2021. A snag survey was not completed in 2026, but we noted that there would likely be a temporary uptick in roost potential due to ash tree mortality as flaking bark would provide temporary roosts, until such time as the bark flaked off.

The acoustic monitor in the FOD5-5 community was set up along the ATV trail on a large shagbark hickory tree from June 18 to June 23. Over that time, a total of 647 bat passes were recorded, of which 6 were from the Little Brown Bat, 33 for the Hoary, 22 for the Eastern Red, and 114 for the Silver Haired. Based on our experience at other mature deciduous forest sites, these numbers are in the low to medium range. None of the SAR bat passes occurred near the

dusk emergence period, suggesting that all were coming in from an adjacent area to forage and were not roosting on site.

In summary, we have few concerns for SAR bat impacts for the following reasons.

1. Much of Flatts Point has limited bat habitat potential.
2. The central area of Flatts Point where development is proposed to be concentrated has minimal bat habitat potential due to the dominance of red cedars.
3. The outer parts of Flatts Point that have better bat potential are not being developed.
4. SAR bat species are not habitat limited in the County.
5. SAR bats are flexible in roost use, changing roosting areas between seasons and within seasons.
6. All bat species are tolerant of nearby human activity, such as roosting in buildings, and we consistently get high bat numbers when recording in urban areas.
7. Bats are only active at night, when most human activity has ended, and therefore the risks of negative bat/human interactions are low.

SAR Bat Recommendations:

The SCA allows the proponent to determine if there is a high level of risk to SAR. In our opinion, Registration with the SCA is not required for SAR bats as long as the following mitigation measures are followed.

1. If there is a need to remove non-red cedar trees greater than 25 cm DBH (measurement assigned by MECP 2022) within the nest/roost season of April 1 to Nov. 30, then the removal of the tree is only recommended if it can be shown that it is not an active roost tree as determined by an acoustic exit survey. Otherwise, there should be no tree clearing of non-red cedar trees greater than 25 cm DBH during the nest/roost season of April 1 to Nov. 30.
2. Unlike the rest of the lots where the first 50 m (in from the lake) of vegetation is recommended to be retained with at least 80% canopy cover, it is recommended that lots 1 and 2 retain the first 100 m. This is not intended to restrict the placement of a well head. If there is a proposal by a future landowner to develop within the 50 to 100 m woodland area of Lots 1 and 2, then it is recommended that development approvals be preceded by SAR bat surveys as per MECP (2022) and the results of these efforts be used to determine if there is a need to Register the development with the SCA. The results of that registration process would have potential impacts on how development would proceed in the 50 to 100 m woodland area.

7.1 Wetland/Coastal Wetland

There is no designated significant wetland within 120 m of the development property. In this regard, we refer to the following excerpt from Section 4.4 of the Natural Heritage Reference Manual.

The need to evaluate the ecological function of adjacent lands (i.e., undertake an EIS or equivalent study) would be removed if proponents choose to avoid having development and site alteration occur within the

extent of adjacent lands.

Accordingly, it is our opinion that no further analysis is warranted for impacts to significant coastal wetland.

There are four small unevaluated coastal wetland patches to the southeast of the development area on Flatts Point. All are separated from Lake Ontario by a cobble beach bar, but all are technically coastal wetlands due to their proximity to the lake. Two are within 120 m, including an approximate 1.5 ha. SWT2 patch, and an approximate 0.1 ha. “w” patch, both previously discussed in the ELC section of this report and shown in Figure 3. From Figure 4, a SWT2-8 patch (~ 0.6 ha) has been identified that will be greater than 120 m from the proposed Lot 17 build location (see Figure 2) and more than 300 m from the eastern boundary of any of the 16 lots on Flatts Point. A patch of MAS2-1 (~0.6 ha) is located between the SWT2-9 and SWT2 patches.

The total combined area of all wetland patches between Lot 17 and the main development is ~2.8 ha., which is slightly above the 2 ha. minimum wetland size that is needed to undertake an evaluation using Ontario Wetland Evaluation System (MNRF 2022). In our opinion the combined patches are too small and lacking in special features to have any potential for provincial significance. We have been undertaking wetland evaluations in the province since 1984 and have completed over 100 full wetland evaluations and many hundreds of wetland assessments since that time and therefore can speak with some authority on this potential significance.

The MNRF (2022) evaluation process scores wetlands on the following four components:

Biological: Primarily intended to score the biodiversity of the wetland. Larger wetlands can do well here as their breadth has a greater probability of having more features to score, and they also automatically receive more points based on size. The wetlands here would receive a low Biological Component score, primarily due to their small size.


Social: Scores wetlands for their potential social value, such as with recreational use. This wetland would get a very low Social Component score due to their small size and isolation on private property.

Hydrological: Scores wetlands primarily for their flood and pollution attenuation abilities. Wetlands on the Great Lakes such as these do not get high Hydrological Component scores because the lake hydrology overrides the wetlands potential value in this regard.

Special Features: Scores wetlands primarily for rare species as well as for unusual biological features and fish habitat. This wetland does not contain fish habitat. A Blanding’s Turtle was observed nearby in 2022 and 2025, suggesting a possible presence in the wetland, but as per MNRF (2022) the score for a single SAR species is not particularly high.

Wetland boundary mapping was based on MNRF (2014). The mapping process is best exemplified by the adjacent flow wetland boundary flow diagram from MNRF (2014). From this, there are three key features that need to be considered: seasonal or permanent water, hydric substrates and coverage by at least 50% wetland vegetation.

Wetland boundary mapping was primarily based on the 50% rule as the wetlands here do not have permanent water, and are primarily located on a firm non-hydric substrate. They are separated from Lake Ontario by a cobble/rock berm varying in height from 1 to 3 meters. Subsurface percolation through the berm from Lake Ontario determines the depth, breadth, and temporal water levels within the wetland as based Lake Ontario water levels. This argument is strengthened by the primarily firm substrates and relatively shallow soils observed in most of the wetland areas. Subsequently, when Lake Ontario goes through its annual seasonal lows (e.g., fall/early winter) there would be no standing water in the wetland. From this we conclude that this wetland exists because of its low topography and the temporary water holding ability of the berm. Relative to this, Lake Ontario water levels have different highs from year to year, and this was observed in the wetland. For example, during low lake level periods, such as during the June 16, 2021 site visit, the wetland had no standing water (see adjacent image). As a result, the wetland would not be able to provide habitat for aquatic dependent species such as certain amphibians and reptiles and fish.





In contrast to the 2021 dry period, there was some ephemeral pooling (~5 cm deep) present (see adjacent) during the June 26, 2020 site and the May 27 and June 7 2026 site visits as the lake was visibly higher at that time. It is our opinion that catchment water flowing from the land side of the wetland does not provide a significant source of water for this wetland, and that it is the lake levels that define water levels in the wetland.

Of the sixteen proposed lots, lots 11 to 16 will be within 120 m of the SWT2 and “w” patch. Due to the depth of these lots it will be possible to push the houses back at least 120 m from the SWT2 and “2” patch, but from the perspective of ameliorating a negative impact to the wetland it is our opinion that a 50 m buffer is sufficient due to a lack of wetland significance and a lack of sensitive features and because SWT2 water levels are controlled by Lake Ontario. As such, the potential for a hydrological impact from the adjacent development is minimal.

Concerns about potential impacts from adjacent septic systems have been raised. Analysis of this type of impact is not the purview of an EIS, but instead will need to be assessed by qualified individuals to ensure compliance with appropriate regulations. We refer to the Hydrological Study (Malroz 2021) and the Terrain Analysis (Groundwork Engineering 2021) that assess the appropriateness of the septic systems and locations.

Aside from that, we do note that the recommended 50 m treed buffer is considered more than sufficient to protect adjacent natural heritage features from pesticides (see Vormeier et al. 2023) and from nutrient impacts (see Moghaddam 2021), with the understanding that 30 m is the standard in the province for such impact purposes, and because treed buffers are considered the more effective in comparison to shrubs and grass (Aguar et al. 2015). In that respect, we do provide the following recommendations for the 50 m buffer.

1. It is recommended that the lots next to wetland areas (Lots 10 to 15 and 17) have an adjacent 50 m no build buffer zone (i.e., residential dwelling, septic fields, sheds). Lot 16 is not next to wetland. It is understood that the well heads will need to be closer to the wetlands and we have no concerns in this regard.
2. It is recommended that shrub and trees next to wetlands in the prescribed 50 m zone of Lots 10 to 15 and 17 that are next to wetland be maintained with at least 80% canopy cover to maintain buffer functionality. The removal of dead or dying trees (such as dead ash trees); non-native and invasive species (e.g., European buckthorn), or trees with a low Coefficient of Conservatism (e.g., Manitoba Maple and Crack Willow) is acceptable if the 80% canopy cover threshold is maintained. If there is a desire to remove these low Coefficient of Conservatism species in the buffer zone below the 80% threshold, they should be replaced by native tree plantings.

7.2 Area of Natural and Scientific Interest (ANSI)

There are no ANSI's within 120 m of the proposed development. In this regard, we refer to the following excerpt from Section 4.4 of the Natural Heritage Reference Manual.

The need to evaluate the ecological function of adjacent lands (i.e., undertake an EIS or equivalent study) would be removed if proponents choose to avoid having development and site alteration occur within the extent of adjacent lands.

Accordingly, it is our opinion that no further analysis is warranted for ANSI's.

7.3 Fish Habitat

Some of the lots will be within 120 m of fish habitat (i.e., Halfmoon Bay). Due to the shallow water, the lack of cover, the lack of aquatic vegetation, and notable wave impacts, we do not consider the offshore fish habitat as important. It also does not contain important fish species as shown on the Department of Fisheries and Oceans Critical Habitat and Distribution Maps.

The nearshore area is comprised of shallow unvegetated flat rock rubble and limestone pavement.



From the Navionics app we see that shallow water (less than 2 m deep) extends about 150 m from shore. The lack of submergent vegetation in this area, which might otherwise support a diverse fish assemblage would be due to a combination of a lack of a rooting substrate, and wave and ice scouring.



The nearshore area is too shallow to provide nesting by non-minnow sized species, and the lack of submergent vegetation and wave action makes it unsuitable for most minnow species nesting except for a few possible lithophilic spawning species such as the longnose dace and darters, although from our experience in similar habitats in Prince Edward County, nearshore nesting would predominately be by the invasive round goby. We do not consider the 6 m zone and beyond as sensitive fish habitat and fish use would mostly be confined to movement and feeding for a low diversity and density of fish.

Salmonids will nest in this kind of substrate, but only if it is next to a deeper shelf, which in most locations is more than 120 m from shore.

For fish habitat we provide the following mitigation recommendations.

1. It is recommended that the lake shoreline areas for lots 1 to 11 (lots 12 to 16 are not next to fish habitat) have a 50 m no build buffer zone (i.e., residential dwelling, septic fields, sheds). It is understood that the well heads will need to be closer to the lake. Due to space constraints within the adjunct lot 17, a 50 m buffer zone to the lake is not feasible, and Regulation 41/24 for development within 30 m will likely need be applied.

2. It is recommended that shrub and trees in the lake buffer zone of Lots 1 to 11 be maintained with at least 80% canopy cover to maintain buffer functionality. The removal of dead or dying trees (e.g., ash); non-native and invasive species (e.g., European buckthorn), or trees with a low Coefficient of Conservatism (e.g., Manitoba Maple and Crack Willow) is acceptable if the 80% canopy cover threshold is maintained. If there is a desire to remove these low Coefficient of Conservatism species in the buffer zone below the 80% threshold, they should be replaced primarily by native tree species. Except for a band of trees along the shoreline that are mostly Crack Willow, the potential build area of Lot 17 is not treed and therefore maintaining an 80% crown cover is not feasible.

7.4 Valleylands

There are no valleylands within 120 m of the proposed development. In this regard, we refer to the following excerpt from Section 4.4 of the Natural Heritage Reference Manual.

The need to evaluate the ecological function of adjacent lands (i.e., undertake an EIS or equivalent study) would be removed if proponents choose to avoid having development and site alteration occur within the extent of adjacent lands.

Accordingly, it is our opinion that no further analysis is warranted for valleyland.

7.5 Woodlands

Woodland significance is based on criteria outlined in the Natural Heritage Reference Manual and outlined in Table 4:

Table 4: Candidate Significant Woodland Threshold Features based on criteria from the 2010 Natural Heritage Reference Manual.			
Significance Category	Significance Threshold	Candidate Habitat Criteria Analysis	Is Threshold Met
Size	County: between 15 and 30% woodland coverage; > 20 ha is significant	Total area including off property woodlands is ~17 ha.	No
Interior Habitat	> 2 ha of interior habitat is significant	Red cedar woodlands do not factor in the calculations (see below), and true core interior habitat is mostly lacking.	No
Proximity (10 ha area threshold)	If woodland is < 30 m of a significant natural heritage features or fish habitat, and it provides some ecological benefit to that adjacent feature	We consider all woodland areas within 30 m of the lake or adjacent wetlands to perform a significant proximity function regardless of size thresholds.	Yes (see below)
Linkages (10 ha area threshold)	If woodland is located between two other significant features	Not located between two significant features	No
Water Protection (5 ha area threshold)	If woodland is located within a sensitive watershed or sensitive groundwater discharge area, sensitive headwater area, watercourse, or fish habitat.	We consider all woodland areas within 30 m of the lake or adjacent wetlands to perform a significant proximity function regardless of size thresholds.	Yes (see below)
Woodland Diversity (10 ha area threshold)	High biodiversity or a declining forest type	Common species found throughout the region.	No
Uncommon Characteristics (5 ha. area threshold)	S1, S2, S3 vegetation community; 8,9,10 coefficient of conservatism (CC); > 10 trees more than 100 years old/ha or > 10 trees > 50 cm DBH or having a basal area of 8m ² for trees greater than 40 cm DBH	No S1,S2,S3 communities and all woodlands have relatively low CC values. The woodland areas where housing/septics are proposed are mostly dominated by red cedar. Historical air imagery indicates that only the FO5-5 woodland has the potential for larger trees and greater age. The FOD5-5 woodland was not assessed for age, DBH, or B.A. characteristics, but since it is below the 5 ha. area sub-threshold, then significance can not be attained.	No
Conclusion and Recommendations: We do not consider the FOC2-1 eastern red cedar woodland an ecological woodland for the sake of woodland core significance ranking because eastern red cedar is considered an invasive in many jurisdictions (Meneguzzo and Liknes (2015) due to its ability to rapidly expand into new areas and suppress the establishment of other habitat types such as deciduous woodlands. It has been our observation through 40 years of surveying these stands in Eastern Ontario and Prince Edward County that they have a low biodiversity and this is also borne out in the literature (e.g., Horncastle et al. (2005), Norris et al. (2001), Briggs et al., (2002), Heinen and O'Connell (2009), Steinaur (2023), and Schnelle (2023)). Their growth pattern is more			

shrub like, and the concept of core habitat for wildlife purposes would not apply to this forest type due to their structural characteristics. This is also indicated by the lack of core habitat songbirds observed in the red cedars during birding efforts.

Proximity/Water Protection:

1. It is recommended that the shoreline/wetland/SWH areas for lots 1 to 15 have a 50 m no build buffer zone (i.e., residential dwelling, septic fields, sheds). It is understood that the well head will need to be closer to the lake. Lot 16 is not adjacent to lake or wetland. Adjunct Lot 17 is adjacent to the lake, but the potential build area is not treed, although it is recommended that shoreline trees be maintained.

2. It is recommended that shrub and trees in the 50 m buffer zone be maintained with at least 80% canopy cover to maintain buffer functionality. The removal of dead or dying trees (e.g., ash); non-native and invasive species (e.g., European buckthorn), or trees with a low Coefficient of Conservatism (e.g., Manitoba Maple and Crack Willow) is acceptable if the 80% canopy cover threshold is maintained. If there is a desire to remove these low Coefficient of Conservatism species in the buffer zone below the 80% threshold, they should be replaced primarily by native tree species.

7.6 Wildlife Habitat

The Significant Wildlife Habitat Criteria for Ecodistrict 6E (MNR 2015b) in Table 5 describes habitat and wildlife requirements and habitat thresholds needed to reach significance.

Table 5: Candidate SWH analysis.			
SWH Type	Candidate Habitat < 120 m	Criteria analysis	Species Thresholds Met
Waterfowl Stopover and Staging Area (Terrestrial)	No	No sheet water fields	No
Waterfowl Stopover and Staging Area (Aquatic)	No	No appropriate open water wetland types within 120 m	No
Shorebird Migratory Stopover Area	Yes	BBS1-2 (Willow Gravel Shrub Beach Type)	No
Raptor Wintering Area	No	No rodent productive fields	No
Bat Hibernacula	No	No caves	No
Bat Maternity Colonies	No	Mature deciduous stands with > 10ha of 25 cm DBH trees not present	No
Turtle Wintering Areas	No	Waters off shore provide poor turtle wintering potential as does the adjacent wetland areas	No
Reptile Hibernacula	No	Lacks physical features that would support overwintering	No
Colonially-Nesting Bird Breeding Habitat (Bank and Cliff)	No	No exposed banks or cliffs	No
Colonially-Nesting Bird Breeding Habitat (Tree/Shrubs)	No	SWD habitat not present	No
Colonially-Nesting Bird Breeding Habitat (Ground)	Yes	Rocky peninsula	No nests observed and NE rock shoal is underwater at certain times of the year
Migratory Butterfly Stopover Area	No	Mostly lacking in fields	No

Landbird Migratory Stopover Area	Yes	FOD woodland near Lake Ontario	No
Deer Yarding Area	No	Known SWH yarding is many km. to the north of Hwy. 401	No
Deer Winter Congregation Area	No	Requires 100 ha. woodlots	No
Rare Vegetation Communities (Alvar, Cliff, Sand Barren, Old Growth Forest, Savannah, Tallgrass Prairie)	No	Site and adjacent natural areas are common habitat types in the region	No
Waterfowl Nesting	Yes	Adjacent SWT type	No
Bald Eagle/Osprey Nesting, Foraging, and Perching	Yes	All woodlands next to lake	No (see Table 8)
Woodland Raptor Nesting	No	No 10 ha of interior woodland habitat < 120 m (also, red cedar woodlands not suitable for nesting)	No
Turtle Nesting	No	No adjacent appropriate ELC types	No
Seeps and Springs	No	Not a headwater area	No
Amphibian Breeding Habitat (Woodland)	No	No ephemeral woodland pools > 500 m ²	No
Amphibian Breeding Habitat (Wetlands)	Yes	Wetlands < 500 m ²	Yes, see Table 6
Woodland Area-Sensitive Bird Breeding Habitat	No	No woodlot > 30 ha. within 120m. Red cedar woodlands of site do not provide habitat for listed species	No
Marsh Bird Breeding Habitat	No	No marsh habitat	No
Open Country Bird Breeding Habitat	No	30 ha. grassland not within 120m	No
Shrub/Early Successional Bird Breeding Habitat	No	10 ha. thickets not within 120m	No
Terrestrial Crayfish	No	N/A to Eastern Ontario	No
Special Concern and Rare Wildlife Species	Yes	S1, S2, S3, and Special Concern Species	Yes, see Table 7a and 7b
Amphibian Movement Corridors	No	Requires significant breeding habitat, and this is not present	No
Deer Movement Corridors	No	No SWH deer wintering	No

Table 6. Marsh Monitoring Protocol results of three site visits (2021).			
Site Visit 2021	Beaufort Scale	Background Noise Code	Combined Call Codes for both sites
May 4 (8 C)	0	0	Chorus Frog (3) Leopard Frog (1)
May 19 (21 C)	0	1	Grey Tree Frog (3)
June 9 (28 C)	0	0	Grey Tree Frog (3)

	Two wetland sites were surveyed for amphibian calling. Site 1 (see w in Figure 3) and Site 2 (see SWT2 in Figure 3). Two species with CC of 3 were recorded and therefore the test for SWH is met. Recommendations: 1. It is recommended that the combined SWT2 and “w” areas have a 50 m no build buffer zone. This is not intended to restrict the placement of the well head. basis. 2. It is recommended that shrub and trees in the 50 m buffer zone be maintained with at least 80% canopy cover to maintain buffer functionality. The removal of dead or dying trees (e.g., ash); non-native and invasive species (e.g., European buckthorn), or trees with a low Coefficient of Conservatism (e.g., Manitoba Maple and Crack Willow) is acceptable if the 80% canopy cover threshold is maintained. If there is a desire to remove these low Coefficient of Conservatism species in the buffer zone below the 80% threshold, they should be replaced primarily by native tree species. Note: In Ontario, Chorus Frogs are classified as Not at Risk. The Western Chorus Frog is listed as Threatened at the federal level, but this designation only applies to federal property. We also note some interesting and recent research by Chen (2025) that has shown genomic symmetry and evolutionary hybridization between the common Boreal and rarer Western Chorus Frogs, that may raise questions about the identification of separate species.
---	--

Rare Species: MNR (2015b) intends S1 to S3 and SC species to apply to Table 7a. Inclusion into Table 7a is based on presence in NHIC Grids 18UP3867, 3866, 3966, 3766, Henson and Brodribb (2005), Ecological Services field work, and other sources (e.g., anecdotal reports, eBirds, etc.).

Table 7a. List of potential rare species that are not endangered or threatened. Species observed are discussed after the table.				
Species	Preferred Habitat	Habitat Present within 120 m	Observed on < 120 m	Impact Concerns
Reptiles				
Snapping Turtle (SC)	Open water wetlands with lily pads.	No	No	No
Milk Snake (SC)	Cosmopolitan with some field preferences	Yes	No	No
Map Turtle (SC)	Riverine/sheltered lake edge	No	No	No
Birds				
Barn Swallow (SC)	Barns	Yes	No	No
Black Tern (SC)	Colonies in shallow marshes, especially in cattails.	No	No	No
Bald Eagle (SC)	Nest in large trees along lakes and large rivers.	Yes	No	No
Eastern Wood Pewee (SC)	Deciduous woodland.	Yes	No	No
Grasshopper Sparrow (SC)	Short grass fields	No	No	No
Upland Sandpiper S2	Fields	Yes	No	No
Wood Thrush (SC)	Deciduous woodland.	Yes	Yes	See below
Whip-poor-will (SC)	Scrub habitat	Yes	No	No
Plants				
American Beachgrass S3	Sand dunes.	No	No	No

Sand Reed Grass S3	Sand dunes.	No	No	No
Rose Swamp Mallow S3 SC	Shoreline marshes.	No	No	No
Bushy Cinquefoil S3	Dune and cobble beach shorelines.	No		No

Table 7b. List of species classified as either Threatened or Endangered until April 2026, under the previous Endangered Species Act. They are listed here with their SRank.				
Species	Preferred Habitat	Habitat Present within 120 m	Observed on < 120 m	Impact Concerns
Bobolink S4B	Hay fields, meadows	Nearby fields > 120 m associated with Smugglers Cover RV Park in 2023	No	No
Cerulean Warbler S3B	Mature Deciduous Woodland	Most records in PEC are either historical or are for migrants	No	No
Chimney Swift S3B	Urban areas with chimneys	None	No	No
Eastern Meadowlark S3	Hay fields, meadows	Nearby fields	Yes in 2021, but not in 2023 and 2026	See below
Least Bittern S4B	Open Water Marsh	Reported in South Bay > 3 km away	No	No
Loggerhead Shrike S2B	Open Alvar	None ... most records in PEC are for migrants	No	No
Red Headed Woodpecker S4B	Parkland	Smugglers Cove RV Park > 120 m	No	No

Wood Thrush: In 2021, three to four (i.e., one of 3 could be a repeat) Wood Thrush were heard calling in the lake edge woodlands on the eastern side of Flatts Point on May 17, and one was heard calling here on May 31 (see Table 8). None were heard here during the June and July site visits of 2020, nor during any of the other visits in 2021, 2023, and 2026. This is a vocal bird that is readily heard throughout the day and that none were heard calling at other times suggests the birds heard calling in 2021 were either migrants passing through, or were transient males trying to attract a female without success. Whether or not breeding was successful, the habitat at the call locations is suitable for breeding.

The North American decline in Wood Thrush numbers is being experienced by many aerial insectivores (Spiller and Dettmers 2019). Suggested reasons for declines have included a deterioration in insect prey caused by pesticides, global climate changes, increased nest predation, nest parasitism, declines in stopover-habitat quality, and especially by changes in habitat quality in their Central America winter range (Taylor (2019), and Taylor and Stutchbury (2016)). The proposed development has no direct association with any of these.

Bird Life International puts the current North American population of Wood Thrush at about 12 million and this bird is well represented in eBird for PEC, and we commonly observe it during our field work throughout Eastern Ontario. The status of this species is being assessed in the current round of the Ontario Bird Breeding Atlas research as it continues to do well in the eastern part of the province (see Torrenta et al. (2022), which is also readily evident in eBird. Although it is sometimes considered a woodland species of larger woodlands, we have observed it in a broad range of habitats including fencerows, small woodland patches,

fragmented woodlands, and urban woodlands, and this has also been documented for Eastern Ontario by Torrenta et al. (2022). The use of urban woodlands by Wood Thrush in Eastern North America is well known, which suggests a tolerance to development, and this was opined by Jirinec (2015) who noted that housing developments are not necessarily detrimental to this species.

There is no lack of habitat in the County and the loss of some woodland at this site would not constitute a negative impact to County Wood Thrush populations, especially as the recommended 50 m wooded buffer along the lake edge where they were observed in May 2021, and 100 m buffer for lots 1 and 2 will provide sufficient nesting habitat for these birds. Finally, Wood Thrush do not favor red cedar woodlands, which are more prevalent in the interior of Flatt's Point where most development will occur.

Wood Thrush Recommendation: As a migratory bird, it is recommended that land clearing occur outside of the breeding bird season (April 1 to Aug. 30) to avoid a contravention of the Migratory Birds Convention Act.

Eastern Meadowlark (THR). A family of Meadowlarks was observed in a meadow/thicket area, south of the development area, on the Half Moon Bay winery property during the 2021 birding visits (see Table 8). All Meadowlark activity was confined to the southeast corner of a ~ 1 ha. area of scrubby field in 2021, surrounded by a larger thicket community. We undertook one Eastern Meadowlark survey in this field in 2023 and two in 2026, and they were not present. From the 2026 site visits it is apparent that there is insufficient field (i.e., > 0.5 ha.) to classify the previous Meadowlark use area as Meadow, and this is reflected in the updated 2026 ELC mapping. In 2021 we classified it as a CUM/CUT (Cultural Meadow/Cultural Thicket) community, and it is now classified as a CUT1 (see Section 6 for details) community making it unsuitable for use by Meadowlarks. Ideal habitat for this bird would be a 5 ha. field (MNRF 2019b), although we have observed them nesting in fields in the 1 ha. size range.

Meadowlark Recommendation: Given the ecological successional changes of the field where the Meadowlark were observed in 2021, and the lack of suitable habitat within the development area, it is our opinion that potential risks to Eastern Meadowlark from the development are low. As with all migratory birds, it is recommended that land clearing occur outside of the breeding bird season (April 1 to Aug. 30) to avoid a contravention of the Migratory Birds Convention Act.

8. References

- Aguiar, T., F. Bortolozzo, F. Hansel, K. Rasera, and M. Ferreira. 2015. Riparian buffer zones as pesticide filters of no-till crops. *Environmental Science and Pollution Research*. 22: 10618-10626.
- Belden & C. 1878. *Illustrated Historical Atlas of the Counties of Hastings and Prince Edward County*.
- Briggs, J., G. Hoch, and L. Johnson. 2002. Assessing the rate, mechanisms, and consequences of the conversion of tallgrass prairie to *Juniperus virginiana* forest. *Ecosystems* 6:578-586.
- Cadman M. and N. Kopysh. 2001. *Ontario Breeding Bird Atlas Guide for Participants*. Bird Studies Canada, Environment Canada: Canadian Wildlife Service, Federation of Ontario Naturalists, OFO, Ontario Ministry of Natural Resources.
- Carter, T., M. Menzel, B. Chapman, and K. Miller. 1999. Summer foraging and roosting behavior of an eastern pipistrelle, *Pipistrellus subflavus*. *Bat Research News* 40:5–6.
- Chen, Y. 2025. Genome evolution across temporal scales, from synteny and evolution spanning >200 million years to genomic divergence and mito-nuclear discordance among congeners since the Miocene. PhD. Thesis. Queen's University.
- Ecological Services. 2001. Life Science Areas of Natural and Scientific (ANSI) in Site District 6E-15. 2001. Prepared for the Ministry of Natural Resources, Kingston Office.
- Environment and Climate Change Canada. 2018. Recovery Strategy for the Little Brown Myotis (*Myotis lucifugus*), the Northern Myotis (*Myotis septentrionalis*), and the Tri-colored Bat (*Perimyotis subflavus*) in Canada. Species at Risk Act Recovery Strategy Series. Environment and Climate Change Canada, Ottawa. ix + 172 pp.
- Environment and Climate Change Canada. 2016. Management Plan for the Snapping Turtle (*Chelydra serpentina*) in Canada [Proposed]. Species at Risk Act Management Plan Series. Ottawa, Environment and Climate Change Canada, Ottawa, iv + 39 p.
- Ernst, C. 1986. Ecology of the Turtle, *Sternotherus odoratus*, in Southeastern Pennsylvania. *Journal of Herpetology* 20: 341-352.
- Gannon, W.L, R.E. Sherwin, and S. Haymond. 2003. On the importance of articulating assumptions when conducting acoustic studies of habitat used by bats. *Wildlife Society Bulletin* 31:45–61.
- Gleason, Henry A. and Arthur Cronquist. 1991. *Manual of the Vascular Plants of Northeastern United States and Adjacent Canada*, Second Edition. D. Van Nostrand, N.Y.
- Gillingham, J., and C. Carpenter. 1978. Snake hibernation: Construction of and observations on a man-made hibernaculum. *Journal of Herpetology* 12: 495-498.

- Groundwork Engineering Limited. 2021. Halfmoon Bay Subdivision Terrain Analysis Report. Project No. GW-21014
- Hartwig, T., and E. Kiviat. 2007. Microhabitat association of Blanding's Turtles in natural and constructed wetlands in Southeastern New York. *Journal of Wildlife Management* 71: 576-582.
- Hayes, J.P. 1997. Temporal variation in activity of bats and the design of echolocation monitoring studies. *Journal of Mammalogy* 78:514–524.
- Heinen, J., and T. O'Connell. 2009. Influence of invasive eastern red cedar on density of breeding warblers in Cross Timbers Forest. *Proceedings of the Fourth International Partners in Flight Conference: Tundra to Tropics* 698-704.
- Henderson, L., and H. Broders. 2008. Movements and resource selection of the Northern Long-Eared Myotis (*Myotis septentrionalis*) in a forest-agricultural landscape. *Journal of Mammalogy*. 89: 952-963.
- Henry, M, D. Thomas, R. Vaudry, and M. Carrier. 2002. Foraging distances and home range of pregnant and lactating little brown bats (*Myotis lucifugus*). *Journal of Mammalogy* 83: 767-774.
- Henson, B.L. and K.E. Brodribb 2005. *Great Lakes Conservation Blueprint for Terrestrial Biodiversity, Volume 2: Ecodistrict Summaries*. Nature Conservancy of Canada.
- Horncastle, V., E. Hellgren, P. Mayer, D. Engle, and D. Leslie. 2004. Differential consumption of Eastern Red Cedar (*Juniperus virginiana*) by avian and mammalian guilds: Implications for tree invasion. *American Midland Naturalist* 152: 255-267.
- Jirinec, V. 2015. Habitat use of the declining Wood Thrush in coastal Virginia. M.Sc. Thesis. Humboldt State University, College of William and Mary.
- Norris, M., J. Blair, L. Johnson, and R. McKane. 2001. Assessing changes in biomass, productivity, and C and N stores following *Juniperus virginiana* forest expansion into tallgrass prairie. *Canadian Journal of Forest Resources* 31:1940–1946.
- Humphrey, C. 2017. Recovery Strategy for the Eastern Small-footed Myotis (*Myotis leibii*) in Ontario. Ontario Recovery Strategy Series. Prepared for the Ontario Ministry of Natural Resources and Forestry, Peterborough, Ontario. vii + 76 pp.
- Keene, B. 2025. Preliminary Stormwater Management Report. Flatt's Point Subdivision Prince Edward County. Jewell Engineering.
- Kurta, A., C. Schumacher, M. Kurta, and S. Demers. 1999. Roost sites of an eastern pipistrelle during late-summer swarming. *Bat Research News* 40:8–9.

- Law, B., J. Anderson, and M. Chidel. 1998. A bat survey in state forests on the south-west slopes region of New South Wales with suggestions of improvements for future surveys. *Australian Zoologist*. 30: 467-479.
- Lee, H., W. Bakowsky, J. Riley, J. Bowles, M. Puddister, P. Uhlig, and S. McMurray. 1998. Ecological Land Classification for Southern Ontario. Natural Heritage Information Centre.
- Malroz Engineering. 2021. Hydrogeological Assessment, Halfmoon Bay, Prince Edward County, Ontario. File: 1437-101.00
- MECP (Ministry of Environment Conservation and Parks). 2019. Clients Guide to Preliminary Screening for Species at Risk.
- MECP, Ministry of Environment Conservation and Parks. 2022. Maternity Roost Surveys (Forests/Woodlands).
- Meeks, R., and G. Ultsch. 1990. Overwintering behavior of Snapping Turtles. *CopeIS* 3:880-884.
- Moghaddam, M. 2021. The effects of riparian buffer strips on the concentration of phosphorous in aquatic systems in relation to land management practices. M.Sc. Thesis. University of Guelph.
- MNR (Ministry of Natural Resources). 2000. Significant Wildlife Habitat Technical Guide. 151 pp. Fish and Wildlife Branch, Technical Section.
- MNR (Ministry of Natural Resources) 2010. Natural Heritage Reference Manual for Natural Heritage Policies of the Provincial Policy Statement, 2005. Second Edition.
- MNR (Ministry of Natural Resources). 2011. Survey Methodology under the Endangered Species Act, 2007: *Dolichonyx oryzivorus* (Bobolink). Ministry of Natural Resources Policy Division Species at Risk Branch.
- MNR (Ministry of Natural Resources). 2011b. Bats and Bat Habitats: Guidelines for Wind Power Projects. Second Edition.
- MNR (Ministry of Natural Resources). 2012. Survey Protocol for Eastern Whip-poor-will (*Caprimulgus vociferous*) in Ontario. Ontario Ministry of Natural Resources, Species at Risk Branch, Peterborough, Ontario. ii + 10 pp.
- MNR (Ministry of Natural Resources). 2013. Reptile and Amphibian Exclusion Fencing: Best Practices, Version 1.0. Species at Risk Branch Technical Note. Prepared for the Ontario Ministry of Natural Resources, Peterborough, Ontario. 11 pp.
- MNRF (Ministry of Natural Resources and Forestry). 2014. Significant Wildlife Habitat Mitigation Support Tool.
- MNRF (Ministry of Natural Resources and Forestry). 2014b. Use of Buildings and Isolated Trees by Species at Risk Bats: Survey Methodology. October 2014.

- MNRF (Ministry of Natural Resources and Forestry) 2015. Technical Note Species at Risk (SAR) Bats. Regional Operations Division.
- MNRF (Ontario Ministry of Natural Resources). 2015b. Significant Wildlife Habitat Ecoregion 6E Criterion Schedule.
- MNRF (Ministry of Natural Resources and Forestry). 2015c. Survey Protocol for Blanding's Turtle (*Emydoidea blandingii*) in Ontario. Species Conservation Policy Branch. Peterborough, Ontario. ii + 16 pp.
- MNRF (Ministry of Natural Resources and Forestry). 2018. Natural heritage information request guide. Regional Operations Division.
- MNRF (Ministry of Natural Resources and Forestry). 2019a. Bobolink General Habitat Description.
- MNRF (Ministry of Natural Resources and Forestry). 2019b. Meadowlark General Habitat Description.
- MNRF (Ministry of Natural Resources and Forestry). 2022. Ontario Wetland Evaluation System. Southern Manual, 4th ed.
- Newton, E.J., and T. Herman. 2009. Habitat, movements, and behaviour of overwintering Blanding's turtles (*Emydoidea blandingii*) in Nova Scotia. Canadian Journal of Zoology 87: 299–309.
- Rowe, J., and E. Moll. 1991. A radiotelemetric study of activity and movements of the Blanding's turtle (*Emydoidea blandingi*) in northeastern Illinois Journal of Herpetology, Vol. 25, No. 2, pp. 178-185.
- Schaefer, K. 2017. Habitat Usage of tri-colored bats (*Perimyotis subflavus*) in western Kentucky and Tennessee post-white nose syndrome. M.Sc. thesis. Murray State University, Kentucky. 26
- Schowalter, D.B., J.R. Gunson, and L.D. Harder. 1979. Life history characteristics of little brown bats (*Myotis lucifugus*) in Alberta. Canadian Field Naturalist. 93: 243-251.
- Schnelle, M. 2023. Eastern Red Cedar: A United States native tree that ranges from useful, to a nuisance, and even invasive in certain environments. Hort Technology. 33: 570-577
- Schummer, M., J. Palframan, E. McNaughton, T. Barney, and S. Petrie. 2012. Comparisons of bird, aquatic macroinvertebrate, and plant communities among dredged ponds and natural wetland habitats at Long Point, Lake Erie, Ontario. Wetlands 32:945-953.
- Sherwin, R., W. Gannon, and S. Haymond. 2000. The efficacy of acoustic techniques to infer differential use of habitat by bats. Acta Chiropterologica 2:145-153.

- Snider, J., and M. Linck. 2011. Habitat use and movement patterns of Blanding's Turtles (*Emydoidea blandingii*) in Minnesota, USA: A landscape approach to species conservation. *Herpetological Conservation and Biology* 7:185-195.
- Spiller, K., and R. Dettmers. 2019. Evidence for multiple drivers of aerial insectivore declines in North America. *The Condor*. 20:1-13.
- Steinauer, R. 2023. Evaluation of native plant communities on Nebraska Game and Parks Commission State Parks. Nebraska Game and Parks Commission -- White Papers, Conference Presentations, & Manuscripts. 82.
- Szuba, K., and B. Naylor. 1998. Forest Raptors and Their Nests in Central Ontario. Southcentral Sciences Section Field Guide FG-03. Ontario Ministry of Natural Resource.
- Taylor, G., and E. Noll. 1989. Movements and hibernation sites of overwintering painted turtles in southern Ontario. *Canadian Journal of Zoology*. 67: 1877-1881.
- Taylor, C. 2019. Effects of natal dispersal and density-dependence on connectivity patterns and population dynamics in a migratory network. *Frontiers in Ecology and Evolution*. 7:354.
- Taylor, C., and B. Stutchbury. 2016. Effects of breeding versus winter habitat loss and fragmentation on the population dynamics of a migratory songbird. *Ecological Applications* 26:424-437.
- Torrenta, R., K. A. Hobson, D. C. Tozer, and M.-A. Villard. 2022. Losing the edge: trends in core versus peripheral populations in a declining migratory songbird. *Avian Conservation and Ecology* 17:28.
- Thomas, D. W. 1988. The distribution of bats in different ages of Douglas-fir forests. *Journal of Wildlife Management* 52:619–626.
- Thorne, T. 2015. The role of islands in the migration of bats across Lake Erie and Lake Ontario: *Lasiurus borealis*, *Lasiurus cinereus*, and *Perimyotis subflavus*. M.Sc. thesis. Western University.
- Ultsch, G. and D. Lee. 1983. Radiotelemetric observations of wintering Snapping Turtles (*Chelydra serpentina*) in Rhode Island. *Journal of the Alabama Academy of Sciences* 54, 200–206
- Ultsch, G. 2006. The ecology of overwintering among turtles: where turtles overwinter and its consequences. *Biological Reviews* 81: 339-367.
- Ultsch, R., and M. Cochran. 1994. Physiology of Northern and Southern Musk Turtles (*Sternotherus odoratus*) during simulated hibernation: *Physiological Zoology* 67: 263-281.
- Valley, R. T., Cross, and P. Radomski. 2004. The role of submersed aquatic vegetation as habitat for fish in Minnesota Lakes, including the implication of non-native plant invasions and

their management. Minnesota Department of Natural Resources Special Publication 160.

Veilleux, J., J. Whitaker, and S. Veilleux. 2003. Tree-roosting ecology of reproductive female eastern pipistrelles, *Pipistrellus subflavus*, in Indiana. *Journal of Mammalogy* 84: 1068-1075.

Vormeier, P., L. Liebmann, O. Weisner, and M. Liess. 2023. Width of vegetated buffer strips to protect aquatic life from pesticide effects. *Water Research*. 231: 119627.

Wolbert, S., A. Zellner, and H. Whidden. 2014. Bat activity, insect biomass, and temperature along an elevational gradient. *Northeastern Naturalist*. 27:72-85.

9. Species Lists

Table 8. Bird species observed during visits and location relative to ELC descriptors from Section 6 of this report. Please note that the June 9 2023 visit was intended to focus on only Special Concern or ES species, and any other bird observed was put down as an incidental observation.								
Species	4/5/21	17/5/21	31/5/21	19/6/21	22/6/21	9/6/23	May 27 & June 7 2026	Area
Alder Flycatcher							x	SWT2
American Crow			3	2	1	x	x	FO-
American Goldfinch		5	3	1	4	x	x	Throughout
American Redstart			2	1	1			FOD, FOMa, FOD5-5
American Robin	1	3	6	1	2	x	x	Throughout
Baltimore Oriole		3	9	2	1	x	x	Throughout
Belted Kingfisher			1	1	3	x	x	BBS1-2
Black-billed Cuckoo			1					Adjacent farmland to SW
Black-and-white Warbler			1	1				FOD7-2, FOD5-5
Black-backed Gull (Greater)				5				NE rock shoal into OAO
Black-backed Gull (Lesser)					1			NE rock shoal into OAO
Black-capped Chickadee		2	2	5	2	x	x	Throughout
Black-throated Green Warbler			1					FOD7-2, likely migrant
Bonaparte's Gull				14	1			NE rock shoal into OAO
Brown-headed Cowbird			1			x	x	CUM/CUT
Blue Jay	2		4	2	2	x	x	Throughout
Brown Thrasher			1			x		FOD7-2
Bufflehead	3							OAO
Canada Goose	2						x	OAO
Caspian Tern	10							NE rock shoal into OAO

Cedar Waxwing			2					FOD7-2
Chestnut Sided Warbler			1		1			FOD7-2, FOD
Chipping Sparrow			1		1			FOD7-2, FOD
Common Grackle			9	2	7	x	x	Throughout
Common Merganser			1					OAo
Common Yellowthroat		1	5	2	2	x	x	Throughout
Common Raven			1					Flyby
Downy Woodpecker		1	1					FOC2-1, FOD7-2
Double-crested Cormorant	1			1	1			OAo, Flyby
Eastern Meadowlark			2		3			CUM/CUT
Eastern Towhee	1	1	1	1	2	x		CUM/CUT, FOMa
European Starling		2						FOD7-2
Field Sparrow		1	1					FOC2-1
Gray Catbird			8	2	1	x	x	Throughout
Green Heron			3					SWT2, Flyby
Great Crested Flycatcher		1	3	1	2	x	x	FO-
Greater Yellowlegs	2							BBS1-2
Herring Gull	6			5	21			NE rock shoal into OAo
Indigo Bunting							x	CUT1
Killdeer		2			4	x		Road, BBS1-2
Least Flycatcher			2					FOD7-2, FOD
Long-tailed Duck	10							OAo
Mallard	2				25			OAo
Merganser								OAo
Mourning Dove		1	3	2	4	x	x	CUM/CUT, SWT2
Nashville Warbler			1					Migrant
Northern Cardinal	1		2	1	1	x	x	Throughout
Northern Flicker		1	4					FOD, FOD7-2
Northern House Wren		1	1					FOD7-2
Red-bellied Woodpecker		1	1					FOD, FOD7-2
Red-eyed Vireo			5	4	2	x	x	FO-
Red-winged Blackbird			12	24	27		x	Throughout
Ring-billed Gull	2			50+	7		x	NE rock shoal into OAo
Rose-breasted Grosbeak		3	1	1				FOD5-5, FOD7-2
Ruffed Grouse			1					FOC2-1
Scarlet Tanager		1		1				FOMa, FOD5-5
Song Sparrow	1	1	5	4	4	x	x	Throughout
Swamp Sparrow			5	1	1		x	SWT2
Tennessee Warbler		1						FOD7-2, Likely migrant
Veery			1					FOD

Warbling Vireo		2	3	2	2			SWT2, FOD, FOMa
White-breasted Nuthatch		2		1				FOD5-5, FOD7-2
White-throated Sparrow			1				x	FOD7-2
Wild Turkey	1							FOC2-1
Wood Thrush		3	1			1		FOMa, FOD5-5, FOD7-2
Yellow Warbler		2	6	5	8		x	Throughout

Table 9. Species list from CUW1-1 community. 44% of plants are non-native (i.e., SNA), which is well above the 20-30% average for natural areas in Ontario (Oldham et al. 1995). The average Coefficient of Conservatism (CC) value is 1.9 on a scale of 0 to 10, indicating a site with low ecological sensitivity. All but one of the native plants have a provincial rank of S5 (very common). The one exception is an S4 (common).

Species	Common Name	S-Rank	CC
<i>Acer platanoides</i>	Norway Maple	SNA	0
<i>Apocynum androsaemifolium</i>	Spreading Dogbane	S5	3
<i>Arctium minus</i>	Common Burdock	SNA	0
<i>Calystegia sepium</i>	Hedge False Bindweed	S5	2
<i>Carex pensylvanica</i>	Pennsylvania Sedge	S5	5
<i>Carya ovata</i>	Shagbark Hickory	S5	6
<i>Dactylis glomerata</i>	Orchard Grass	SNA	0
<i>Daucus carota</i>	Wild Carrot	SNA	0
<i>Desmodium canadense</i>	Canada Tick-trefoil	S4	5
<i>Eleocharis acicularis</i>	Needle Spikerush	S5	5
<i>Galium palustre</i>	Common Marsh Bedstraw	SNA	0
<i>Geum fragarioides</i>	Barren Strawberry	S5	5
<i>Hydrophyllum virginianum</i>	Virginia Waterleaf	S5	6
<i>Ilex verticillata</i>	Common Winterberry	S5	5
<i>Iris versicolor</i>	Harlequin Blue Flag	S5	5
<i>Juniperus virginiana</i>	Eastern Red Cedar	S5	4
<i>Leucanthemum vulgare</i>	Oxeye Daisy	SNA	0
<i>Lotus corniculatus</i>	Garden Bird's-foot Trefoil	SNA	0
<i>Medicago lupulina</i>	Black Medick	SNA	0
<i>Melilotus albus</i>	White Sweet-clover	SNA	0
<i>Phalaris arundinacea</i>	Reed Canarygrass	S5	0
<i>Phleum pratense</i>	Common Timothy	SNA	0
<i>Pilosella piloselloides ssp. praealta</i>	King Devil Hawkweed	SNA	0
<i>Poa pratensis</i>	Kentucky blue grass	SNA	0
<i>Potentilla norvegica</i>	Rough Cinquefoil	S5	0
<i>Ranunculus abortivus</i>	Kidney-leaved Buttercup	S5	2
<i>Rhus aromatica</i>	Fragrant Sumac	S4	8
<i>Rhamnus cathartica</i>	European Buckthorn	SNA	0
<i>Rhus typhina</i>	Staghorn Sumac	S5	1
<i>Rosa blanda</i>	Smooth Rose	S5	3
<i>Solidago canadensis</i>	Canada Goldenrod	S5	1
<i>Solidago juncea</i>	Early Goldenrod	S5	3

<i>Sonchus arvensis</i>	Field Sow-thistle	SNA	0
<i>Stellaria longifolia</i>	Long-leaved Starwort	S5	2
<i>Symphyotrichum novae-angliae</i>	New England Aster	S5	2
<i>Taraxacum officinale</i>	Common Dandelion	SNA	0
<i>Tilia americana</i>	Basswood	S5	4
<i>Tragopogon dubius</i>	Yellow Goatsbeard	SNA	0
<i>Trifolium hybridum</i>	Alsike Clover	SNA	0
<i>Ulmus americana</i>	White Elm	S5	3
<i>Urtica dioica</i> ssp. <i>dioica</i>	European Stinging Nettle	SNA	0
<i>Vincetoxicum rossicum</i>	Dog Strangling Vine	SNA	0
<i>Xanthium strumarium</i>	Rough Cocklebur	S5	2

Table 10. Species list from CUT1 community. 46 % of plants are non-native (i.e., SNA), which is above the 20-30% average for natural areas in Ontario (Oldham et al. 1995). The average Coefficient of Conservatism (CC) value is 1.8 on a scale of 0 to 10, indicating a site with low ecological sensitivity. All but one of the native plants have a provincial rank of S5 (very common). The one exception is an S4 (common).

Species	Common Name	S-Rank	CC
<i>Acer platanoides</i>	Norway Maple	SNA	0
<i>Apocynum androsaemifolium</i>	Spreading Dogbane	S5	3
<i>Arctium minus</i>	Common Burdock	SNA	0
<i>Bromus inermis</i>	Smooth Brome	SNA	0
<i>Calystegia sepium</i>	Hedge False Bindweed	S5	2
<i>Carex pensylvanica</i>	Pennsylvania Sedge	S5	5
<i>Dipsacus fullonum</i>	Teasel	SNA	0
<i>Leonurus cardiaca</i>	Motherwort	SNA	0
<i>Conyza canadensis</i>	Horseweed	SNA	0
<i>Carya ovata</i>	Shagbark Hickory	S5	6
<i>Cornus racemosa</i>	Gray dogwood	S5	2
<i>Dactylis glomerata</i>	Orchard Grass	SNA	0
<i>Daucus carota</i>	Wild Carrot	SNA	0
<i>Desmodium canadense</i>	Canada Tick-trefoil	S4	5
<i>Eleocharis acicularis</i>	Needle Spikerush	S5	5
<i>Galium palustre</i>	Common Marsh Bedstraw	S5	5
<i>Geum fragarioides</i>	Barren Strawberry	S5	5
<i>Hydrophyllum virginianum</i>	Virginia Waterleaf	S5	6
<i>Ilex verticillata</i>	Common Winterberry	S5	5
<i>Iris versicolor</i>	Harlequin Blue Flag	S5	5
<i>Juniperus virginiana</i>	Eastern Red Cedar	S5	4
<i>Leucanthemum vulgare</i>	Oxeye Daisy	SNA	0
<i>Lotus corniculatus</i>	Garden Bird's-foot Trefoil	SNA	0
<i>Medicago lupulina</i>	Black Medick	SNA	0
<i>Melilotus albus</i>	White Sweet-clover	SNA	0
<i>Phalaris arundinacea</i>	Canary reed grass	SNA	0
<i>Phleum pratense</i>	Common Timothy	SNA	0
<i>Pilosella piloselloides</i> ssp. <i>praealta</i>	King Devil Hawkweed	SNA	0
<i>Potentilla anserina</i>	Silverweed	S5	5
<i>Potentilla norvegica</i>	Rough Cinquefoil	S5	0
<i>Ranunculus abortivus</i>	Kidney-leaved Buttercup	S5	2
<i>Rhamnus cathartica</i>	European Buckthorn	SNA	0

<i>Rhus typhina</i>	Staghorn Sumac	S5	1
<i>Rosa blanda</i>	Smooth Rose	S5	3
<i>Rubus idaeus</i>	Wild red raspberry	SNA	0
<i>Rubus occidentalis</i>	Black raspberry	S5	2
<i>Solidago canadensis</i>	Canada Goldenrod	S5	1
<i>Solidago juncea</i>	Early Goldenrod	S5	3
<i>Sonchus arvensis</i>	Field Sow-thistle	SNA	0
<i>Stellaria longifolia</i>	Long-leaved Starwort	S5	2
<i>Symphyotrichum novae-angliae</i>	New England Aster	S5	2
<i>Taraxacum officinale</i>	Common Dandelion	SNA	0
<i>Tilia americana</i>	Basswood	S5	4
<i>Tragopogon dubius</i>	Yellow Goatsbeard	SNA	0
<i>Trifolium hybridum</i>	Alsike Clover	SNA	0
<i>Ulmus americana</i>	White Elm	S5	3
<i>Urtica dioica</i> ssp. <i>dioica</i>	European Stinging Nettle	SNA	0
<i>Xanthium strumarium</i>	Rough Cocklebur	S5	2

Table 11. Species list from the FOD5-5 woodland. 12% of plants are non-native (i.e., SNA), which is below the 20-30% average for natural areas in Ontario (Oldham et al. 1995). The average Coefficient of Conservatism (CC) value is 3.1 on a scale of 0 to 10, indicating a site with low ecological sensitivity. All but one of the native plants have a provincial rank of S5 (very common). The one exception is an S4 (common).

Species	Common Name	S-Rank	CC
<i>Acer negundo</i>	Manitoba Maple	S5	0
<i>Acer platanoides</i>	Norway Maple	SNA	0
<i>Acer rubrum</i>	Red Maple	S5	4
<i>Acer saccharum</i>	Sugar Maple	S5	4
<i>Achillea millefolium</i>	Common Yarrow	SNA	0
<i>Alliaria petiolata</i>	Garlic Mustard	SNA	0
<i>Anemonastrum canadense</i>	Canada Anemone	S5	3
<i>Carex pensylvanica</i>	Pennsylvania Sedge	S5	5
<i>Carex rosea</i>	Rosy Sedge	S5	2
<i>Carya cordiformis</i>	Bitternut Hickory	S5	6
<i>Carya ovata</i>	Shagbark Hickory	S5	6
<i>Desmodium canadense</i>	Canada Tick-trefoil	S5	5
<i>Fragaria virginiana</i>	Wild Strawberry	S5	2
<i>Fraxinus americana</i>	White Ash	S4	4
<i>Fraxinus pennsylvanica</i>	Green Ash	S4	3
<i>Geranium robertianum</i>	Herb-Robert	S5	2
<i>Geum fragarioides</i>	Barren Strawberry	S5	5
<i>Geum laciniatum</i>	Rough Avens	S4	4
<i>Hydrophyllum virginianum</i>	Virginia Waterleaf	S5	6
<i>Juglans nigra</i>	Black Walnut	S5	5
<i>Ostrya virginiana</i>	Eastern Hop-hornbeam	S5	4
<i>Parthenocissus vitacea</i>	Thicket Creeper	S5	4
<i>Podophyllum peltatum</i>	May-apple	S5	5
<i>Populus tremuloides</i>	Trembling aspen	S5	2
<i>Populus balsamifera</i>	Balsam poplar	S5	4
<i>Populus deltoides</i>	Eastern Cottonwood	S5	4

<i>Prunella vulgaris</i>	Common Self-heal	S5	0
<i>Prunus serotina</i>	Black Cherry	S5	3
<i>Prunus virginiana</i>	Chokecherry	S5	2
<i>Quercus rubra</i>	Northern Red Oak	S5	6
<i>Ranunculus abortivus</i>	Kidney-leaved Buttercup	S5	2
<i>Rhamnus cathartica</i>	European Buckthorn	SNA	0
<i>Ribes cynosbati</i>	Eastern Prickly Gooseberry	S5	4
<i>Solidago flexicaulis</i>	Zigzag Goldenrod	S5	6
<i>Symphyotrichum cordifolium</i>	Heart-leaved Aster	S5	5
<i>Thuja occidentalis</i>	Eastern White Cedar	S5	4
<i>Tilia americana</i>	Basswood	S5	4
<i>Toxicodendron radicans</i>	Poison Ivy	S5	2
<i>Ulmus americana</i>	White Elm	S5	3
<i>Vincetoxicum rossicum</i>	Dog Strangling Vine	SNA	0
<i>Vitis riparia</i>	Riverbank Grape	S5	0

Table 12. Species list from the FOD7 woodland. 22% of plants are non-native (i.e., SNA), which is within the 20-30% average for natural areas in Ontario (Oldham et al. 1995). The average Coefficient of Conservatism (CC) value is 2.6 on a scale of 0 to 10, indicating a site with low ecological sensitivity. All but one of the native plants have a provincial rank of S5 (very common). The one exception is an S4 (common).

Species	Common Name	S-Rank	CC
<i>Acer negundo</i>	Manitoba Maple	S5	0
<i>Acer rubrum</i>	Red Maple	S5	4
<i>Acer saccharum</i>	Sugar Maple	S5	4
<i>Achillea millefolium</i>	Common Yarrow	SNA	0
<i>Alliaria petiolata</i>	Garlic Mustard	SNA	0
<i>Anemonastrum canadense</i>	Canada Anemone	S5	3
<i>Carex pensylvanica</i>	Pennsylvania Sedge	S5	5
<i>Carex rosea</i>	Rosy Sedge	S5	2
<i>Carya cordiformis</i>	Bitternut Hickory	S5	6
<i>Fragaria virginiana</i>	Wild Strawberry	S5	2
<i>Fraxinus pennsylvanica</i>	Green Ash	S4	3
<i>Parthenocissus vitacea</i>	Thicket Creeper	S5	4
<i>Prunus virginiana</i>	Chokecherry	S5	2
<i>Ranunculus abortivus</i>	Kidney-leaved Buttercup	S5	2
<i>Rhamnus cathartica</i>	European Buckthorn	SNA	0
<i>Symphyotrichum cordifolium</i>	Heart-leaved Aster	S5	5
<i>Toxicodendron radicans</i>	Poison Ivy	S5	2
<i>Ulmus americana</i>	White Elm	S5	3
<i>Acer saccharinum</i>	Silver Maple	S5	5
<i>Ambrosia artemisiifolia</i>	Common Ragweed	S5	0
<i>Carex lupulina</i>	Hop Sedge	S5	6
<i>Carex stipata</i>	Awl-fruited Sedge	S5	3
<i>Circaea alpina</i>	Small Enchanter's Nightshade	S5	6
<i>Glyceria striata</i>	Fowl Mannagrass	S5	3
<i>Hesperis matronalis</i>	Dame's Rocket	SNA	0
<i>Onoclea sensibilis</i>	Sensitive Fern	S5	4
<i>Persicaria maculosa</i>	Spotted Lady's-thumb	SNA	0
<i>Rhamnus cathartica</i>	European Buckthorn	SNA	0

<i>Sambucus canadensis</i>	Common Elderberry	S5	5
<i>Urtica dioica</i> ssp. <i>dioica</i>	European Stinging Nettle	SNA	0
<i>Zanthoxylum americanum</i>	Prickly ash	S5	3

Table 13. Species list from the FOM woodland. 26% of plants are non-native (i.e., SNA), which is within the 20-30% average for natural areas in Ontario (Oldham et al. 1995). The average Coefficient of Conservatism (CC) value is 2.4 on a scale of 0 to 10, indicating a site with low ecological sensitivity. All but one of the native plants have a provincial rank of S5 (very common). The one exception is an S4 (common).

Species	Common Name	S-Rank	CC
<i>Acer negundo</i>	Manitoba Maple	S5	0
<i>Acer saccharum</i>	Sugar Maple	S5	4
<i>Alliaria petiolata</i>	Garlic Mustard	SNA	0
<i>Carex pensylvanica</i>	Pennsylvania Sedge	S5	5
<i>Carex rosea</i>	Rosy Sedge	S5	2
<i>Carya cordiformis</i>	Bitternut Hickory	S5	6
<i>Fraxinus pennsylvanica</i>	Green Ash	S4	3
<i>Rhamnus cathartica</i>	European Buckthorn	SNA	0
<i>Symphotrichum cordifolium</i>	Heart-leaved Aster	S5	5
<i>Ulmus americana</i>	White Elm	S5	3
<i>Glyceria striata</i>	Fowl Mannagrass	S5	3
<i>Hesperis matronalis</i>	Dame's Rocket	SNA	0
<i>Sambucus canadensis</i>	Common Elderberry	S5	5
<i>Fraxinus americana</i>	White Ash	S4	4
<i>Geranium robertianum</i>	Herb-Robert	S5	2
<i>Ostrya virginiana</i>	Eastern Hop-hornbeam	S5	4
<i>Ribes cynosbati</i>	Eastern Prickly Gooseberry	S5	4
<i>Solidago flexicaulis</i>	Zigzag Goldenrod	S5	6
<i>Thuja occidentalis</i>	Eastern White Cedar	S5	4
<i>Tilia americana</i>	Basswood	S5	4
<i>Vincetoxicum rossicum</i>	European Swallowwort	SNA	0
<i>Vitis riparia</i>	Riverbank Grape	S5	0
<i>Juniperus communis</i>	Common Juniper	S5	4
<i>Juniperus virginiana</i>	Eastern Red Cedar	S5	4
<i>Lonicera tatarica</i>	Tatarian Honeysuckle	SNA	0
<i>Malus pumila</i>	Common Apple	SNA	0
<i>Medicago lupulina</i>	Black Medick	SNA	0
<i>Poa pratensis</i>	Kentucky Bluegrass	S5	0
<i>Populus deltoides</i>	Eastern Cottonwood	S5	4
<i>Populus tremuloides</i>	Trembling Aspen	S5	2
<i>Ranunculus acris</i>	Common Buttercup	SNA	0
<i>Rubus idaeus</i>	Red Raspberry	S5	2
<i>Rubus occidentalis</i>	Black Raspberry	S5	2
<i>Vincetoxicum rossicum</i>	Dog Strangling Vine	SNA	0
<i>Zanthoxylum americanum</i>	Common Prickly-ash	S5	3

Table 14. Species list from the FOC2-1 woodland. 52% of plants are non-native (i.e., SNA), which is well above the 20-30% average for natural areas in Ontario (Oldham et al. 1995). The average Coefficient of Conservatism (CC) value is 1.3 on a scale of 0 to 10, indicating a site with low

ecological sensitivity. All but one of the native plants have a provincial rank of S5 (very common). The one exception is an S4 (common).

Species	Common Name	S-Rank	CC
<i>Achillea millefolium</i>	Common Yarrow	SNA	0
<i>Ambrosia artemisiifolia</i>	Common Ragweed	S5	0
<i>Arctium minus</i>	Common Burdock	SNA	0
<i>Bromus inermis</i>	Smooth Brome	SNA	0
<i>Carex rosea</i>	Rosy Sedge	S5	2
<i>Dactylis glomerata</i>	Orchard Grass	SNA	0
<i>Daucus carota</i>	Wild Carrot	SNA	0
<i>Fraxinus pennsylvanica</i>	Green Ash	S4	3
<i>Geum fragarioides</i>	Barren Strawberry	S5	5
<i>Juniperus virginiana</i>	Eastern Red Cedar	S5	4
<i>Leucanthemum vulgare</i>	Oxeye Daisy	SNA	0
<i>Parthenocissus vitacea</i>	Thicket Creeper	S5	4
<i>Persicaria maculosa</i>	Spotted Lady's-thumb	SNA	0
<i>Potentilla argentea</i>	Silvery Cinquefoil	SNA	0
<i>Ranunculus acris</i>	Common Buttercup	SNA	0
<i>Rhus typhina</i>	Staghorn Sumac	S5	1
<i>Rhamnus cathartica</i>	European Buckthorn	SNA	0
<i>Ribes cynosbati</i>	Eastern Prickly Gooseberry	S5	4
<i>Thuja occidentalis</i>	Eastern White Cedar	S5	4
<i>Tilia americana</i>	Basswood	S5	4
<i>Urtica dioica</i> ssp. <i>dioica</i>	European Stinging Nettle	SNA	0
<i>Vincetoxicum rossicum</i>	Dog Strangling Vine	SNA	0
<i>Xanthium strumarium</i>	Rough Cocklebur	S5	2
<i>Zanthoxylum americanum</i>	Common Prickly-ash	S5	3

10. Qualifications

Dale Kristensen (botany)

Education

- Masters of Science: Biology Queen's University, Kingston, Ontario, 1996.
- Bachelor of Science: Biology University of Guelph, Guelph Ontario. 1981.

Affiliations and Certifications

- Ecological Restoration Society
- North American Wildflower Society
- Land Conservancy for Kingston, Frontenac, Lennox & Addington
- Kingston Field Naturalists
- COSEWIC Species Recovery Team – Deerberry and Cerulean Warbler habitat modelling.

Positions

1988 – present: Environmental Consultant and co-founder of Ecological Services, Elginburg, Ontario.

1985-2019 Program Associate, Phytotron Manager, and Curator of the Fowler Herbarium. Department of Biology at Queen's University, Kingston, Ontario.

Rob Snetsinger (all taxa)

Education

- Masters of Science (1993): Biology Queen's University, Kingston, Ontario.
- Bachelor of Science (1985): Biology Queen's University, Kingston, Ontario.
- Diploma (1978): Forestry Lakehead University, Thunder Bay, Ontario.

Certifications

- Ontario Wetland Evaluation Assessor
- Ecological Land Classification
- Butternut Health Assessor: Ontario (#123)
- NHIC Data Sensitivity Training

Positions

1984 – present: Environmental Consultant. Co-founder of Ecological Services, Elginburg, Ontario.
1985-2018 Program Associate. Department of Biology at Queen's University, Kingston, Ontario.

Kurt Hennige (breeding bird surveys)

Education

Mechanical Engineering Technician Degree, Polytechnical Institute, Würzburg, Germany

Affiliations

- Lennox & Addington Stewardship Council, Past-Chair 2011- 2018
- Kingston Field Naturalists, President 2014 - 2016
- Birding organizer with Kingston Field Naturalists, Lennox and Addington Stewardship Council, Migration Research Foundation, MNR and Bird Studies Canada
- Breeding Bird Surveys (for Canadian Wildlife Service) 1991- 2025
- E-Bird reviewer for Lennox and Addington, Frontenac, and Leeds and Grenville counties.

Positions

- Avifauna Surveyor, Ecological Services
- Nature Conservancy Canada, Whip-poor-will and Loggerhead Shrike outreach
- Shrike Biologist and Stewardship Habitat Coordinator, Ontario Ministry of Natural Resources and Wildlife Preservation Canada
- Marsh Monitoring, Quinte Conservation
- Bobolink and Eastern Meadowlark Project Lead, Lennox and Addington Stewardship Council
- Species at Risk Surveyor, Nature Conservancy of Canada
- Species at Risk Surveyor, Blue Earth Renewables
- Birding Tour Guide, guiding multiple birding trips around the world and in Canada

Kaitlyn Closs (amphibian survey)

Education

Diploma (2023). Conservation Biology. Flemings College School of Environmental and Natural Resource Sciences.

Training

Trained by Ecological Services in 2022 to undertake snake, turtle, and amphibian surveys.

Positions

2022 – 2025: Ecological surveys for Ecological Services, Elginburg, Ontario.

2022 – 2025: White Lake Provincial Park. A variety of roles since 2022, and is currently the supervisor of outdoor operations.

Jo Barr

Ms. Barr has been trained by Ecological Services to complete turtle, amphibian, and evening avifauna surveys, for which she has been completing on multiple projects since 2024. When needed, she also acts as a field assistant.